

# SYMMETRIC FUNCTIONS AND MACDONALD POLYNOMIALS

ROBIN LANGER

Submitted in total fulfilment of the requirements of the degree of  
Master of Mathematics by Research at the University of Melbourne

ABSTRACT. The ring of symmetric functions  $\Lambda$ , with natural basis given by the Schur functions, arise in many different areas of mathematics. For example, as the cohomology ring of the grassmanian, and as the representation ring of the symmetric group. One may define a coproduct on  $\Lambda$  by the plethystic addition on alphabets. In this way the ring of symmetric functions becomes a Hopf algebra. The Littlewood–Richardson numbers may be viewed as the structure constants for the co-product in the Schur basis. The first part of this thesis, inspired by the umbral calculus of Gian-Carlo Rota, is a study of the co-algebra maps of  $\Lambda$ . The Macdonald polynomials are a somewhat mysterious  $qt$ -deformation of the Schur functions. The second part of this thesis contains a proof a generating function identity for the Macdonald polynomials which was originally conjectured by Kawanaka.

**Declaration.** *The first part of this thesis is entirely the author's own work. The idea for the second part was suggested by Ole Warnaar who also carried out the derivation of equation 15 on pages 59–60 and verified proposition 2.1. The proof of lemma 2.1 was suggested by Paul Zinn-Justin. Steps two and three in the proof of the second part of the thesis are the author's own work.*

**Acknowledgements.** *The author would like to thank Ole Warnaar for suggesting the subject of Macdonald polynomials, Peter Forrester, Alain Lascoux, Paul Zinn-Justin and the two anonymous referees for useful comments and suggestions on the manuscript as well as the University of Montreal for hosting an interesting lecture series and workshop on Macdonald Polynomials and Combinatorial Hopf Algebras.*

*for Mrs McLaren*

## CONTENTS

|                                                             |    |
|-------------------------------------------------------------|----|
| 1. Symmetric functions of Littlewood–Richardson type        | 7  |
| 1.1. Symmetric Functions                                    | 9  |
| 1.1.1. Partitions                                           | 9  |
| 1.1.2. Monomial symmetric functions                         | 15 |
| 1.1.3. Plethystic notation                                  | 16 |
| 1.1.4. Schur functions                                      | 17 |
| 1.2. The Umbral Calculus                                    | 18 |
| 1.2.1. Coalgebras                                           | 18 |
| 1.2.2. Sequences of Binomial Type                           | 22 |
| 1.3. The Hall inner-product                                 | 26 |
| 1.3.1. Preliminaries                                        | 26 |
| 1.3.2. Column operators                                     | 28 |
| 1.3.3. Duality                                              | 30 |
| 1.4. Littlewood–Richardson Bases                            | 33 |
| 1.4.1. Generalized complete symmetric functions             | 34 |
| 1.4.2. Umbral operators                                     | 36 |
| 1.4.3. Column operators                                     | 37 |
| 1.4.4. Generalized elementary symmetric functions           | 40 |
| 1.5. Examples                                               | 41 |
| 2. A generating function identity for Macdonald polynomials | 47 |
| 2.1. Macdonald Polynomials                                  | 47 |
| 2.1.1. Notation                                             | 47 |
| 2.1.2. Operator definition                                  | 48 |
| 2.1.3. Characterization using the inner product             | 49 |
| 2.1.4. Arms and legs                                        | 49 |
| 2.1.5. Duality                                              | 50 |
| 2.1.6. Kawanaka conjecture                                  | 50 |
| 2.2. Resultants                                             | 52 |
| 2.2.1. Residue calculations                                 | 53 |
| 2.3. Pieri formula and recurrence                           | 56 |
| 2.3.1. Arms and legs again                                  | 57 |
| 2.3.2. Pieri formula                                        | 58 |
| 2.3.3. Recurrence                                           | 59 |
| 2.4. The Proof                                              | 59 |
| 2.4.1. The Schur case                                       | 59 |
| 2.4.2. Step one                                             | 62 |
| 2.4.3. Step two                                             | 65 |
| 2.4.4. Step three                                           | 70 |
| References                                                  | 75 |

## 1. SYMMETRIC FUNCTIONS OF LITTLEWOOD–RICHARDSON TYPE

The Schur polynomials  $\{s_\lambda(X)\}$  arise in both geometry and representation theory as a natural basis for the ring of symmetric polynomials which we denote by  $\Lambda$ . The Littlewood–Richardson numbers  $\{c_{\mu\nu}^\lambda\}$  are defined by:

$$s_\mu(X)s_\nu(X) = \sum_{\lambda} c_{\mu\nu}^\lambda s_\lambda(X)$$

The plethystic addition of alphabets gives a coproduct structure on  $\Lambda$ , which induces, via the Hall inner product, the natural product structure on the dual space  $\Lambda^*$  - which may be thought of as the ring of symmetric “formal power series” in some dual alphabet  $Y$ .

Since the Schur polynomials are self-dual with respect to the Hall inner product, the Littlewood–Richardson numbers may also be defined by:

$$s_\lambda(X + X') = \sum_{\mu, \nu} c_{\mu\nu}^\lambda s_\mu(X)s_\nu(X')$$

A sequence of binomial type is a basis  $p_n(x)$  for the one variable polynomial ring  $\mathbb{Q}(x)$  with the property that:

$$p_n(x + x') = \sum_{k=0}^n \binom{n}{k} p_k(x)p_{n-k}(x')$$

Sequences of binomial type arise as images, under co-algebra maps, of the standard basis  $\{x^n\}$ . Furthermore, each such co-algebra map, or *umbral operator* is associated to an a formal power series  $f(z)$  which is invertible with respect to function composition.

The homomorphism from the one variable polynomial ring  $\mathbb{Q}[x]$  to the ring of symmetric functions  $\Lambda$  given by:

$$\frac{x^n}{n!} \mapsto h_n(X)$$

preserves the coproduct structure. Similarly, the homomorphism from the ring of formal power series in one variable  $\mathbb{Q}[[y]]$  to  $\Lambda^*$  given by:

$$y^n \mapsto m_{(n)}(Y)$$

preserves the product structure.

In the special case where the alphabet  $X$  contains a single variable, both these maps are in fact isomorphisms, and the Hall inner product becomes the bracket used in the umbral calculus of Gian-Carlo Rota:

$$\langle f(y), p(x) \rangle = \mathcal{L}f(\mathcal{D})[p(x)]$$

Here  $\mathcal{L}$  denotes the constant term operator, while  $\mathcal{D}$  is the differential operator.

The Schur functions may be expressed as either a determinant in the complete symmetric functions:

$$s_\lambda(X) = \det(h_{\lambda_i+j-i}(X))$$

or as a determinant in the elementary symmetric functions:

$$s_\lambda(X) = \det(e_{\lambda'_i+j-i}(X))$$

The main result of the first part of this thesis is the following. Let  $\{p_n(x)\}$  be the sequence of binomial type arising as the image of  $\{x^n\}$  under the umbral operator  $U_f$  which is associated to the invertible formal power series  $f(z)$ . Let  $\{r_n(X)\}$  be the image of  $\{\frac{p_n(x)}{n!}\}$  under the embedding of  $\mathbb{Q}[x]$  into  $\Lambda$  just described. Then the vector space basis for  $\Lambda$  defined by:

$$P_\lambda(X) = \det(r_{\lambda_i+j-i}(X))$$

has the Littlewood–Richardson property:

$$P_\lambda(X + X') = \sum_{\mu, \nu} c_{\mu\nu}^\lambda P_\mu(X) P_\nu(X')$$

Furthermore, suppose that  $g(z)$  is the compositional inverse of  $f(z)$  and that  $\{q_n(x)\}$  is the sequence of binomial type arising as the image of the standard basis  $\{x^n\}$  under the umbral operator  $U_g$ . Let  $\{\tilde{r}_n(X)\}$  denote the image of  $\{\frac{q_n(x)}{n!}\}$  under the embedding of  $\mathbb{Q}[x]$  into  $\Lambda$  and let:

$$c_n(X) = \omega[\tilde{r}_n(X)]$$

where  $\omega$  is the natural involution on  $\Lambda$  which maps the complete symmetric functions to the elementary symmetric functions.

The aforementioned basis  $\{P_\lambda(X)\}$  has an alternate expression of the form:

$$P_\lambda(X) = \det(c_{\lambda_i+j-i}(X))$$

Every basis  $\{P_\lambda(X)\}$  which has the Littlewood–Richardson property, and which also admits an expansion of the form:

$$P_\lambda(X) = \sum_{\mu \subseteq \lambda} \square_{\lambda\mu} s_\mu(X)$$

arises in this way.

In section 1.1 we review some classical results from the theory of symmetric functions. In section 1.2 we review classical results from the umbral calculus, in section 1.3 we prove the main theorem and in section 1.4 we give some examples.



**1.1. Symmetric Functions.** The main references for this section are Macdonald [29] and Lascoux [24]. Another good reference is Stanley [42]. Also, see Bergeron's online notes (in French) [2]. For connections with the geometry of grassmanians see Fulton [8] and Manivel [30]. For connections with the representation theory of the symmetric and general linear groups see Fulton [8] and Sagan [41]. For more about  $\lambda$ -rings see Knutson [22]. For more about the  $q$ -exponential functions and  $q$ -binomial coefficients see the book by Kac and Cheung [20] or the online notes by Foata and Han [7]. The standard reference for basic hypergeometric series is Gasper and Rahman [14]. For an interesting approach to obtaining basic hypergeometric series identities by specialization of symmetric functions see Bowman [3]

**1.1.1. Partitions.** For our purposes, a composition is just a list of non-negative integers, finitely many of which are greater than zero. For most purposes we will identify lists of different lengths which differ only by a tail of zeros.

A *partition* is a composition whose parts are weakly decreasing. If the sum of the parts of  $\lambda$  is  $n$  then we say that  $\lambda$  is a partition of  $n$  and write  $\lambda \vdash n$ .

Some authors do not allow partitions to contain zero parts, however there are some circumstances in which it is convenient to envision the partition as sitting inside some rectangle of pre-determined dimensions, in which case the tail of zeros becomes significant.

The set of all partitions of  $n$  with  $k$  parts (any number of which may be zero) is denoted by  $\mathcal{P}_k^n$ . The generating series for partitions is the first  $q$ -exponential function:


$$E_q(z) = \frac{1}{(z; q)_\infty} = \sum_{n,k} |\mathcal{P}_k^n| q^n z^k = \sum_k \frac{z^k}{(q; q)_k}$$

Here we are making use of the hypergeometric notation:

$$(a; q)_n = \prod_{k=0}^{n-1} (1 - aq^k)$$

It is common practice to represent a partition pictorially as rows of boxes. The *conjugate* of a partition is obtained by reflecting in the main diagonal. The prime symbol is used to indicate conjugation. Conjugating a partition interchanges the role of rows and columns. The parts of  $\lambda'$  correspond to the columns of  $\lambda$ .

Below, on the left, is the diagram for the partition  $\lambda = (4, 2, 1)$  and on the right, its conjugate  $\lambda' = (3, 2, 1, 1)$



We are using the English convention.

We define the length of a partition, denoted by  $l(\lambda)$ , to be the number of nonzero parts. With this convention, the length of a partition is always less than or equal to its number of parts. For example,  $\lambda = (3, 2, 2, 1, 0, 0)$  is a partition of 8 of length 4 with 6 parts.

There are two natural additions on the set of partitions. The first one corresponds to the concatenation of the individual parts:

$$\lambda + \mu = (\lambda_1 + \mu_1, \lambda_2 + \mu_2, \dots, \lambda_n + \mu_n)$$

The second corresponds to taking the union of the two sets of parts, and then re-ordering them as appropriate. For example:

$$(5, 3, 1) \cup (7, 3, 2, 2) = (7, 5, 3, 3, 2, 2, 1)$$

One may check that:

$$\lambda' + \mu' = (\lambda \cup \mu)'$$

Of particular interest are partitions that contain no repeated parts, including no repeated zero parts. The smallest such partition with  $k$  distinct parts is the staircase partition:

$$\delta_k = (k-1, k-2, \dots, 2, 1, 0)$$

Observe that  $\delta_k$  is a partition of  $k(k-1)/2$ . If  $\lambda$  is a partition of  $n$  into  $k$  parts then  $\lambda + \delta_k$  is a partition of  $n + k(k-1)/2$  into  $k$  distinct parts. Conversely, every partition of  $n + k(k-1)/2$  into  $k$  distinct parts is equal to  $\lambda + \delta_k$  for some  $\lambda \vdash n$ .

We denote by  $\mathcal{D}_k^n$  the set of all partitions of  $n + k(k-1)/2$  into  $k$  distinct parts (one of which may be zero). The generating series for partitions with distinct parts is the second  $q$ -exponential function:

$$e_q(z) = (-z; q)_\infty = \sum_{n,k} |\mathcal{D}_k^n| q^n z^k = \sum_k q^{\binom{k}{2}} \frac{z^k}{(q; q)_k}$$

Note that the conjugate of a partition with distinct parts will not, in general, be another partition with distinct parts. In fact, the only

partition with distinct parts whose conjugate also has distinct parts is the staircase partition  $\delta_k$ .

It is perhaps worth pointing out here, though we won't need it until much later, that if  $\pi_q$  is the function which maps  $q$  to  $1/q$  then:

$$\pi_q[(q; q)_n] = (-1)^n q^{-\frac{n(n+1)}{2}} (q; q)_n$$

and so:


$$\pi_q[E_q(z)] = e_q(-qz)$$

If the diagram for a partition  $\mu$  sits properly inside the diagram for the partition  $\lambda$ , that is, if each part of  $\mu$  is less than or equal to the corresponding part of  $\lambda$  then we write  $\mu \subseteq \lambda$ . For such a pair of partitions, we may form the *skew partition*  $\lambda/\mu$  which is simply the collection of boxes in  $\lambda$  which are not also boxes of  $\mu$ .

A skew partition  $\lambda/\mu$  is said to be a *horizontal strip* if:

$$\lambda_1 \geq \mu_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq \mu_n$$

That is, no column contains more than a single box.




For an arbitrary partition  $\mu$  we let  $U(\mu)$  denote the set of partitions  $\lambda$  such that  $\lambda/\mu$  is a horizontal strip, and let  $D(\mu)$  denote the set of partitions  $\nu$  such that  $\mu/\nu$  is a horizontal strip.

More specifically, we let  $U_r(\mu)$  denote the set of partitions  $\lambda$  such that  $\lambda/\mu$  is a horizontal strip with exactly  $r$  boxes, and let  $D_r(\mu)$  denote the set of partitions  $\nu$  such that  $\mu/\nu$  is a horizontal strip with exactly  $r$  boxes.

Similarly a skew partition  $\lambda/\mu$  is a *vertical strip* if:

$$\lambda'_1 \geq \mu'_1 \geq \lambda'_2 \geq \dots \geq \lambda'_n \geq \mu'_n$$

That is, no row contains more than a single box.



More specifically, we let  $\tilde{U}_r(\mu)$  denote the set of partitions  $\lambda$  such that  $\lambda/\mu$  is a vertical strip with exactly  $r$  boxes, and let  $\tilde{D}_r(\mu)$  denote the set of partitions  $\nu$  such that  $\mu/\nu$  is a vertical strip with exactly  $r$  boxes.

An  $(n, m)$ -binomial path is a binary string containing exactly  $n$  zeros and  $m$  ones. Pictorially, we may represent an  $(n, m)$ -binomial path by drawing an  $n$  by  $m$  grid and tracing a path from the bottom left hand corner to the top right hand corner by reading the binary string from left to right and taking a step up each time we read a one, and a step across each time we read a zero.

[illegible]

$\alpha(s) =$ 

|  |  |  |  |
|--|--|--|--|
|  |  |  |  |
|  |  |  |  |
|  |  |  |  |
|  |  |  |  |

$\beta(s) =$ 

|  |  |  |  |
|--|--|--|--|
|  |  |  |  |
|  |  |  |  |
|  |  |  |  |
|  |  |  |  |


$$\tilde{\mathcal{S}} =$$

It is not hard to see that the upper diagram associated with a binomial path  $s$  is equal to the lower diagram associated with the dual

path  $\tilde{s}$ . Likewise the lower diagram of  $s$  is equal to the upper diagram of  $\tilde{s}$ .

$$\alpha(\tilde{s}) = \begin{array}{|c|c|c|c|c|} \hline & & & & \\ \hline & & & & \\ \hline & & & & \\ \hline & & & & \\ \hline \end{array} = \beta(s) \quad \beta(\tilde{s}) = \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array} = \alpha(s)$$


Now let  $s'$  denote the binary string obtained from  $s$  by swapping all the zeros for ones and vice versa. In our case  $s' = 1100101011$ . Note that  $s'$  is now an  $(m, n)$  binomial path.



Observe that  $(\alpha(s'), \beta(s')) = (\beta(s)', \alpha(s'))$ , and similarly  $(\alpha(\tilde{s}'), \beta(\tilde{s}')) = (\beta(\tilde{s})', \alpha(\tilde{s})') = (\alpha(s)', \beta(s'))$ . In other words, reversing the string and interchanging the role of zeros and ones is equivalent to conjugating the partition.

$$\alpha(s') = \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array} = \beta(s)' \quad \beta(s') = \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array} = \alpha(s)'$$

There is also a way that we may associate a pair of diagrams  $(\nu(s), \gamma(s))$  with *distinct* parts to any given binomial path. Suppose that we label the steps of  $s$  from left to right with the integers from 0 to  $n + m - 1$ .



Then  $\nu(s)$  is the diagram which has a row of length  $k$  if and only if there is some upstep of  $s$  labelled with  $k$ . We have of course that  $\nu(s) = \alpha(s) + \delta_n$

The other partition  $\gamma(s)$  is the diagram which has a row of length  $k$  if and only if there is some *across* step which is labelled with a  $k$ .

Of course  $\nu(s) \cup \gamma(s) = \delta_{m+n}$ .

Performing the same procedure with the dual path  $s'$  we get:

This time:

and:

Observe that, a little surprisingly, we have  $\gamma(s) = \nu(s')$  and  $\nu(s) = \gamma(s')$ , which gives us the following proposition:

**Proposition 1.1.** *Let  $\lambda$  be any partition contained within an  $n$  by  $m$  box, and let  $\mu$  be its compliment with respect to this box, then we have:*

$$(\lambda + \delta_n) \cup (\mu' + \delta_m) = \delta_{n+m}$$

*Proof.* Suppose that  $\lambda = \alpha(s)$ , so that  $\lambda + \delta_n = \nu(s)$ . Now  $\mu' = \alpha(s')$  and  $\mu' + \delta_m = \nu(s') = \gamma(s)$ . The result now follows from the fact that  $\nu(s) \cup \gamma(s) = \delta_{n+m}$ .  $\square$

1.1.2. *Monomial symmetric functions.* Consider now the multivariate polynomial ring  $\mathbb{Q}[x_1, \dots, x_k]$ . Each monomial in  $\mathbb{Q}[x_1, \dots, x_k]$  corresponds to a composition. For example, the monomial  $x_1 x_3^2$  in  $\mathbb{Q}[x_1, x_2, x_3]$  corresponds to the list  $(1, 0, 2)$ .

For notational convenience, if  $\eta = (\eta_1, \eta_2, \dots, \eta_k)$  is a composition, then by  $X^\eta$  we mean the monomial  $x_1^{\eta_1} x_2^{\eta_2} \dots x_k^{\eta_k}$ . There is a natural addition on the space of compositions which corresponds to multiplication in the polynomial ring. if  $\eta = (\eta_1, \dots, \eta_k)$  and  $\gamma = (\gamma_1, \dots, \gamma_k)$  then  $\eta + \gamma = (\eta_1 + \gamma_1, \dots, \eta_k + \gamma_k)$ , and  $X^{(\eta+\gamma)} = X^\eta X^\gamma$ .

The symmetric group  $\mathcal{S}_k$  acts on the set of compositions with  $k$  parts by permuting the parts. Each orbit of  $\mathcal{S}_k$  contains a unique partition. For any composition  $\lambda$  let  $r(\lambda, i)$  denote the number of parts of  $\lambda$  equal to  $i$ . Let us define:

$$r_\lambda = \prod_{i \geq 0} r(\lambda, i)!$$

Then  $r_\lambda$  is the order of the subgroup of  $\mathcal{S}_k$  that stabilizes the composition  $\lambda$ . If  $\lambda$  is a partition with distinct parts then  $r_\lambda = 1$ .

For each partition  $\lambda$  we define the *monomial symmetric polynomial* to be:

$$m_\lambda = \frac{1}{r_\lambda} \sum_{\sigma \in S_n} X^{\sigma(\lambda)}$$

Since  $r_\lambda$  is the number of permutations in  $\mathcal{S}_k$  that will stabilize the monomial  $x^\lambda$ , the monomial symmetric function is the sum of all *distinct* permutations of this monomial.

As  $\lambda$  runs over the set of all partitions of  $n$  with at most  $k$  parts, the set  $\{m_\lambda | \lambda \in \mathcal{P}_n^k\}$  forms a basis for the vector space  $\Lambda_n^k$  consisting of all symmetric polynomials of homogeneous degree  $k$  in  $n$  variables. We shall write:

$$\Lambda_n = \bigoplus_k \Lambda_n^k$$

to denote the ring of symmetric functions in  $n$  variables. The map:

$$\pi_n : \Lambda_n \rightarrow \Lambda_{n-1}$$

which sets the  $n$ th variable equal to zero is a homomorphism whose kernel is generated by the monomial symmetric functions indexed by partitions with exactly  $n$  nonzero parts.

1.1.3. *Plethystic notation.* It is often more convenient to work in infinitely many variables. It is also convenient to make use of the plethystic notation. In the plethystic notation one writes:  $X = x_1 + x_2 + x_3 + \cdots$  to denote the set of indeterminates:  $X = \{x_1, x_2, x_3, \dots\}$  and  $Y = y_1 + y_2 + y_3 + \cdots$  to denote the set of indeterminates  $Y = \{y_1, y_2, y_3, \dots\}$

Furthermore by  $X + Y$  we denote the disjoint union of the variables in the sets  $X$  and  $Y$

$$\begin{aligned} X + Y &= x_1 + x_2 + \cdots + y_1 + y_2 + \cdots \\ &= \{x_1, x_2, \dots, y_1, y_2, \dots\} \\ &= X \cup Y \end{aligned}$$

and by  $XY$  we denote the cartesian product:

$$\begin{aligned} XY &= (x_1 + x_2 + \cdots)(y_1 + y_2 + \cdots) \\ &= x_1y_1 + x_1y_2 + \cdots + x_2y_1 + x_2y_2 + \cdots \\ &= \{x_1y_1, x_1y_2, \dots, x_2y_1, x_2y_2, \dots\} \\ &= X \times Y \end{aligned}$$

The complete and elementary symmetric functions may be defined in terms of their generating functions:

$$\begin{aligned} \Omega_z(X) &= \Omega(Xz) = \prod_{x \in X} \frac{1}{1 - xz} = \sum_{n=0}^{\infty} h_n(X) z^n \\ \tilde{\Omega}_z(X) &= \tilde{\Omega}(Xz) = \prod_{x \in X} (1 + xz) = \sum_{n=0}^{\infty} e_n(X) z^n \end{aligned}$$

Note that the first  $q$ -exponential function may be expressed as:

$$E_q(z) = \Omega\left(\frac{z}{1-q}\right)$$

where  $\frac{1}{1-q}$  is the alphabet  $1 + q + q^2 + \dots$ . Similarly the second  $q$ -exponential function may be expressed as:

$$e_q(z) = \tilde{\Omega}\left(\frac{z}{1-q}\right)$$

The involution  $\omega$  is defined by:

$$\omega[h_n(X)] = e_n(X)$$



(extended multiplicatively). The “forgotten” symmetric functions are defined by:

$$\omega[m_\lambda(X)] = f_\lambda(X)$$

For each variable  $x$  let us define an *anti-variable*  $\bar{x}$  such that:

$$\{x\} \cup \{\bar{x}\} = \{\}$$

Alternatively one can think in terms of *anti-sets*:

$$\{x\} \cup \overline{\{x\}} = \{\}$$

Now let  $-X = \bar{x}_1 + \bar{x}_2 + \bar{x}_3 + \cdots$  denote the set of anti-variables  $\{\bar{x}_1, \bar{x}_2, \bar{x}_3, \dots\}$  or, the *anti-set* of variables  $\overline{\{x_1, x_2, x_3, \dots\}}$ .

For an arbitrary symmetric function  $f$  we define:

$$f(-X) = \omega f(\epsilon X)$$

where  $\epsilon X$  is the more obvious but less natural form of negation:

$$\epsilon X = \{-x_1, -x_2, -x_3, \dots\}$$

We may now express the elementary symmetric functions in terms of the complete symmetric functions:

$$\tilde{\Omega}(X) = \Omega(-\epsilon X)$$

1.1.4. *Schur functions.* The most important basis for the ring of symmetric functions, from the perspective of geometry or representation theory, is the Schur functions.

Let  $H(X)$  denote the infinite Toeplitz matrix  $(h_{i-j}(X))$ , where we are assuming that  $h_{-k}(X) = 0$  for all  $k \geq 1$ . In other words the matrix is upper triangular with ones on the diagonal. For  $\lambda$  and  $\mu$  partitions with  $n$  parts (any number of which may be zero), the skew Schur function  $s_{\lambda/\mu}(X)$  may be defined by minors of this matrix as follows:

$$s_{\lambda/\mu}(X) = \det(H_{I,J}(X)) = \det(h_{\lambda_i - \mu_j + j - i}(X))$$

where  $I = \mu + \delta_n$  and  $J = \lambda + \delta_n$ . Note that we indexing both the rows and the columns from zero rather than one. In particular the regular Schur functions are given by:

$$s_\lambda(X) = \det(H_{[n], \lambda + \delta_n}) = \det(h_{\lambda_i + j - i}(X))$$

where  $[n] = \{0, 1, 2, \dots, n-1\}$ . Note that  $n$  does not appear explicitly in this equation, and if  $n$  is taken to be larger than the number of nonzero parts of  $\lambda$  then the resulting matrix is block diagonal, with the first block of size  $l(\lambda)$  independent of  $n$ , and the remaining blocks containing ones, which do not affect the determinant.

The fact that the generating function for the complete symmetric functions has the property  $\Omega_z(X + X') = \Omega_z(X)\Omega_z(X')$  lifts to the fact that:  $H(X + X') = H(X)H(X')$ , and so, by the Cauchy–Binet theorem we have:

$$s_\lambda(X + X') = \sum_{\mu \subseteq \lambda} s_{\lambda/\mu}(X) s_\mu(X')$$

Consider now the infinite matrix  $E(X) = (e_{i-j}(X))$ . The fact that  $\Omega(X)\tilde{\Omega}(\epsilon X) = 1$  now lifts to the fact that  $H(X)E(\epsilon X) = I$ . Thus by Jacobi's formula for the minors of the inverse matrix, we have:

$$\begin{aligned} \det(H_{I,J}(X)) &= (-1)^{\sum_{k=1}^r j_k - i_k} \det(E_{J',I'}(\epsilon X)) \\ &= \det(E_{J',I'}(X)) \end{aligned}$$

Now by proposition 1.1, if these are  $(n + m)$  by  $(n + m)$  matrices and  $I = \lambda + \delta_n$  then  $I' = \lambda' + \delta_m$ . In other words we have an alternative expression for the Schur functions given by:

$$s_{\lambda/\mu}(X) = \det(e_{\lambda'_i - \mu'_j + j - i}(X))$$

We also get from this that:

$$\omega[s_\lambda(X)] = s_{\lambda'}(X)$$

**1.2. The Umbral Calculus.** A nice reference for finite dimensional linear algebra is Hoffman and Kunze [17]. For more about algebras, co-algebras, Hopf algebras and quantum groups see the online notes by Arun Ram [36]. The book by Roman [38] discusses the infinite-dimensional subtleties more carefully than is done here.

**1.2.1. Coalgebras.** An algebra is a vector space  $V$  over some field  $F$  equipped with a multiplication:

$$m : V \otimes V \rightarrow V$$

and a unit:

$$e : F \rightarrow V$$

such that the following two diagrams commute:

$$\begin{array}{ccc} V \otimes V \otimes V & \xrightarrow{m \otimes id} & V \otimes V \\ \downarrow id \otimes m & & \downarrow m \\ V \otimes V & \xrightarrow{m} & V \end{array}$$

This is just the associative law.

$$\begin{array}{ccc}
 V & \xrightarrow{e \otimes id} & V \otimes V \\
 \downarrow id \otimes e & \searrow id & \downarrow m \\
 V \otimes V & \xrightarrow{m} & V
 \end{array}$$

This says that multiplication by the identity has no effect.

A commutative algebra is an algebra with the additional property that:

$$\begin{array}{ccc}
 V \otimes V & \xrightarrow{\tau} & V \otimes V \\
 \searrow m & & \swarrow m \\
 & V &
 \end{array}$$

where  $\tau : V \otimes V \rightarrow V \otimes V$  is the map  $\tau(x \otimes y) = y \otimes x$ .

The ring  $\Lambda$  of symmetric functions, and the ring  $\mathbb{Q}[x]$  of polynomials in one variable are both commutative algebras.

An algebra morphism is a vector space morphism:  $\psi : V \rightarrow V$  satisfying the two properties:

$$\begin{array}{ccc}
 V \otimes V & \xrightarrow{\psi \otimes \psi} & W \otimes W \\
 \downarrow m & & \downarrow m \\
 V & \xrightarrow{\psi} & W
 \end{array}$$

The multiplication is preserved.

$$\begin{array}{ccc}
 V & \xrightarrow{\psi} & W \\
 \swarrow e & & \searrow e \\
 & F &
 \end{array}$$

The identity is preserved.

The map which sends the complete symmetric function  $h_n(X)$  to  $x^n$  is an algebra morphism from  $\Lambda$  onto  $\mathbb{Q}[x]$ .

A coalgebra is a vector space  $V$  equipped with a comultiplication:

$$\Delta : V \rightarrow V \otimes V$$

and a counit:

$$c : V \rightarrow F$$

such that the following diagrams commute:

$$\begin{array}{ccc} V \otimes V \otimes V & \xleftarrow{\Delta \otimes id} & V \otimes V \\ id \otimes \Delta \uparrow & & \uparrow \Delta \\ V \otimes V & \xleftarrow{\Delta} & V \end{array}$$

This is the co-associative law.

$$\begin{array}{ccc} V & \xleftarrow{c \otimes id} & V \otimes V \\ id \otimes c \uparrow & \nearrow id & \uparrow \Delta \\ V \otimes V & \xleftarrow{\Delta} & V \end{array}$$

This says that co-multiplying by the co-unit has no affect.

A co-commutative co-algebra is a co-algebra with the additional property that:

$$\begin{array}{ccc} V \otimes V & \xleftarrow{\tau} & V \otimes V \\ & \searrow \triangleright & \swarrow \triangleright \\ & V & \end{array}$$

Both the ring of symmetric functions and the ring of polynomials are co-commutative co-algebras. In the case of  $\mathbb{Q}[x]$  we have:

$$\mathbb{Q}[x] \otimes \mathbb{Q}[x] \cong \mathbb{Q}[x, x']$$

and the coproduct is given by:

$$\Delta[p(x)] = p(x + x')$$

In the symmetric function case we have, when working in an infinite alphabet, that:

$$\Lambda \otimes \Lambda \cong \Lambda$$

The coproduct is given by the plethystic addition of alphabets.

A co-algebra morphism is a vector space morphism:  $\phi : V \rightarrow V$  satisfying the two properties:

$$\begin{array}{ccc} V \otimes V & \xleftarrow{\phi \otimes \phi} & W \otimes W \\ \Delta \uparrow & & \uparrow \Delta \\ V & \xleftarrow{\phi} & W \end{array}$$

The comultiplication is preserved.

$$\begin{array}{ccc} & \phi & \\ V & \xleftarrow{\quad} & W \\ & \searrow \epsilon & \swarrow \epsilon \\ & F & \end{array}$$

The counit is preserved.

The map which sends  $x^n/n!$  to the complete symmetric function  $h_n(X)$  is a co-algebra morphism from  $\mathbb{Q}[x]$  into  $\Lambda$ . We shall make use of this homomorphism to lift properties of  $\mathbb{Q}[x]$  to  $\Lambda$ .

Categorically, the notions of algebra and co-algebra are dual. Combinatorially the product describes how the elements of an algebra can be combined, while the coproduct describes how the elements of a co-algebra may be decomposed [19, 39].

Every vector space  $V$  has a dual  $V^*$  which is the space of linear functionals from  $V$  into the field  $F$ . There is a natural map:

$$(-, -) : V^* \otimes V \rightarrow F$$

given by:

$$(w, v) = w[v]$$

In finitely many dimensions, every vector space is isomorphic to its dual, and once an isomorphism  $\phi : V \rightarrow V^*$  has been fixed we may define a non-degenerate bilinear form:

$$\langle -, - \rangle : V \otimes V \rightarrow F$$

by:

$$\langle v_1, v_2 \rangle = (\phi(v_1), v_2)$$

Still in finite dimensions, if  $V$  is an algebra then  $V^*$  is a coalgebra with coproduct:

$$(\Delta w, v_1 \otimes v_2) = (w, m(v_1, v_2))$$

and counit:

$$c(w) = (w, e(1))$$

Furthermore, if  $\psi : V \rightarrow W$  is an algebra morphism, then its adjoint  $\psi^* : W^* \rightarrow V^*$  is a coalgebra morphism.

Similarly, if  $V$  is a co-algebra then  $V^*$  is an algebra with multiplication:

$$(m(w_1, w_2), v) = (w_1 \otimes w_2, \Delta(v))$$

and unit:

$$(e(x), v) = xc(v)$$

If  $\phi : V \rightarrow V$  is an co-algebra morphism, then its adjoint  $\phi^* : V^* \rightarrow V^*$  is an algebra morphism.

Unfortunately, both  $\Lambda$  and  $\mathbb{Q}[x]$  are infinite dimensional. In infinitely many dimensions, a vector space  $V$  need not be isomorphic to its dual  $V^*$ , and instead of equality we have:

$$V^* \otimes V^* \subseteq (V \otimes V)^*$$

Since  $\Delta(w) \in V^* \otimes V^*$  whenever  $w \in V^*$  we are still able, in infinite dimensions, to induce a product structure on the dual from an existing product structure on the original space, and we still have that algebra morphisms of  $V^*$  correspond to co-algebra morphisms of  $V$ . The converse is however no longer true.

**1.2.2. Sequences of Binomial Type.** The classification of the co-algebra isomorphisms of the polynomial ring  $\mathbb{Q}[x]$  by consideration of the algebra morphisms of the dual space goes back to Gian-Carlo Rota [33, 40]. Since then there have been many other nice expositions such as Gessel [15] and Roman [38], as well as attempts to find  $q$ -analogs [16, 18, 37], and attempts to generalize the umbral calculus to symmetric functions: Loeb [26], Mendez [31, 32] and Chen [4, 5]. We follow here, in particular, the presentation of Garsia [9, 11, 12]. See the following online resources [6, 25] for a more extensive bibliography.

The ring of polynomials in one indeterminate  $V = \mathbb{Q}[x]$  has a distinguished Hamel basis given by:

$$\{1, x, \frac{x^2}{2!}, \frac{x^3}{3!}, \dots\}$$

The dual space  $V^* = \mathbb{Q}[[y]]$  has a distinguished Schauder basis:

$$\{1, y, y^2, y^3, \dots\}$$

We may think of  $y^k$  as the  $k$ th coefficient extraction operator (acting on exponential generating series):

$$(y^k, p(x)) = \left[ \frac{x^k}{k!} \right] p(x)$$

Note that coefficient extraction and differentiation are related by:

$$\left[ \frac{x^k}{k!} \right] f(x) = \mathcal{L} \mathcal{D}^k f(x)$$

where  $\mathcal{D}$  denotes the ordinary differential operator  $d/dx$ , and  $\mathcal{L}$  is the constant term operator. Thus we may also write the pairing between  $V$  and  $V^*$  as:

$$(f(y), p(x)) = \mathcal{L} f(\mathcal{D})[p(x)]$$

By the Leibniz rule, the co-multiplication on  $V$ :

$$\Delta p(x) = p(x + x')$$

induces the multiplication on  $V^*$ :

$$m(y^r, y^s) = y^{r+s}$$

Define a delta series to be a formal power series  $f(y)$  which has a compositional inverse. For every delta series  $f(y)$  the map:

$$\phi_f(y) = f(y)$$

induces an algebra morphism of  $V^*$  with inverse:

$$\phi_g(y) = g(y)$$

where  $g(y)$  is the compositional inverse of  $f(y)$ , that is:

$$f(g(y)) = g(f(y)) = y$$

A sequence of polynomials  $\{p_n(x)\}$  is said to be of *binomial type* if:

$$p_n(x + x') = \sum_k \binom{n}{k} p_k(x) p_{n-k}(x')$$

A sequence of polynomials  $\{p_n(x)\}$  is said to be of *convolution type* if:

$$p_n(x + x') = \sum_k p_k(x) p_{n-k}(x')$$

Clearly, if  $p_n(x)$  is of binomial type, then  $p_n(x)/n!$  is of convolution type.

Sequences of convolution type arise as images, under co-algebra maps, of the standard basis  $\{x^n/n!\}$  while the corresponding sequences of binomial type arise as images, under co-algebra maps, of the basis  $\{x^n\}$ .

By duality, the co-algebra maps of  $V$  are each adjoint to an algebra map of  $V^*$ . We shall write  $U_f = \phi_f^*$  to denote the umbral operator defined by:

$$U_f \left[ \frac{x^n}{n!} \right] = p_n(x)$$

where  $\{p_n(x)\}$  is the convolution type basis of  $V$  which is dual to the basis  $\{1, g(y), g(y)^2, \dots\}$  of  $V^*$ . As before  $g(y)$  is the compositional inverse of  $f(y)$ .

Let  $f_1(y) = \sum_k a_k y^k$  and  $f_2(y) = \sum_k b_k y^k$  be delta series. Now let:

$$\begin{aligned} \alpha_{nk} &= [y^n] f_2(y)^k \\ \beta_{nk} &= [y^n] f_1(y)^k \\ \gamma_{nk} &= [y^n] f_1(f_2(y))^k \end{aligned}$$

and let  $A = (\alpha_{nk})$  and  $B = (\beta_{nk})$  and  $C = (\gamma_{nk})$  be infinite matrices with  $n, k \geq 1$ . Since:

$$\begin{aligned} f_1(f_2(y))^n &= \sum_i \beta_{in} f_2(y)^i \\ &= \sum_k \left( \sum_i \alpha_{ki} \beta_{in} \right) y^k \end{aligned}$$

We have that:

$$AB = C$$

These are called Jabotinsky matrices [12].

**Proposition 1.2.** *The umbral operator has the explicit expression:*

$$U_f = \sum_n \frac{x^n}{n!} \mathcal{L} f^n(\mathcal{D})$$

*Alternatively, the umbral operator may be characterized by the two properties:*

$$(i) \quad \mathcal{D} \circ U_f = U_f \circ g(\mathcal{D})$$

$$(ii) \quad \mathcal{L} \circ U_f = \mathcal{L}$$



*Proof.* For the first part, note that:

$$[y^n]f(y) = f(\mathcal{D}) \left[ \frac{x^n}{n!} \right]$$

Let  $A = \{a_{nk}\}$  be the matrix:

$$a_{nk} = [y^k]f(y)^n$$

and let  $B = \{b_{nk}\}$  be the inverse matrix:

$$b_{nk} = [y^k]g(y)^n$$

We have:

$$\begin{aligned} \langle g(y)^k, U_f[x^n/n!] \rangle &= \mathcal{L}g(\mathcal{D})^k \sum_j \frac{x^j}{j!} [y^n]f(y)^j \\ &= \sum_j [y^j]g(y)^k [y^n]f(y)^j \\ &= \sum_j b_{kj} a_{jn} \\ &= \delta_{nk} \end{aligned}$$

For the second part, we have:

$$\begin{aligned} \langle g(y)^k, g(\mathcal{D})[p_n(x)] \rangle &= \mathcal{L}g(\mathcal{D})^{k+1}[p_n(x)] \\ &= \delta_{k+1,n} \\ &= \delta_{k,n-1} \\ &= \langle g(y)^k, p_{n-1}(x) \rangle \end{aligned}$$

Which shows that the umbral operator satisfies the first property. The second property is obvious. To see that these properties characterize the umbral operator, recall Taylor's formula:

$$\mathcal{I} = \sum_n \frac{x^n}{n!} \mathcal{L}\mathcal{D}^n$$

Suppose that  $\mathcal{U}$  is any operator satisfying the above two properties, then we have:

$$\begin{aligned}
 \mathcal{U} &= \sum_n \frac{x^n}{n!} \mathcal{L} \mathcal{D}^n \mathcal{U} \\
 &= \sum_n \frac{x^n}{n!} \mathcal{L} \mathcal{U} \circ f^n(\mathcal{D}) \\
 &= \sum_n \frac{x^n}{n!} \mathcal{L} \circ f^n(\mathcal{D}) \\
 &= \mathcal{U}_f
 \end{aligned}$$

□

**1.3. The Hall inner-product.** The results in this section are taken from the book of Lascoux [24] and the two papers by Garsia, Haiman and Tesler [10, 13]. See also the paper by Zabrocki [43].

1.3.1. *Preliminaries.* The Hall inner product may be defined by:

$$\langle m_\lambda(Y), h_\mu(X) \rangle = \delta_{\lambda\mu}$$

Normally this is thought of as a map:

$$\langle -, - \rangle : \Lambda \otimes \Lambda \rightarrow \mathbb{Q}$$

but we want to think of it as a map:

$$\langle -, - \rangle : \Lambda^* \otimes \Lambda \rightarrow \mathbb{Q}$$

In analogy with the one variable case,  $\Lambda$  is symmetric polynomials while  $\Lambda^*$  is symmetric “formal power series”. In lieu of the fact that a pair of bases  $\{P_\lambda(X)\}$  and  $\{Q_\lambda(Y)\}$  are dual with respect to the Hall inner product if and only if:

$$\sum_\lambda P_\lambda(X) Q_\lambda(Y) = \Omega(XY) = \prod_{\substack{x \in X \\ y \in Y}} \frac{1}{1 - xy}$$

We shall always write elements of  $\Lambda$  as symmetric functions in the alphabet  $X$  and elements of  $\Lambda^*$  as symmetric functions in the alphabet  $Y$ .

Any symmetric function  $f(X)$  may be thought of as the operator “multiplication by  $f(X)$ ”, and as such there is an adjoint operator, which we denote by  $\partial_f$ . By definition:

$$\langle \partial_f[g(Y)], h(X) \rangle = \langle g(Y), f(X)h(X) \rangle$$

The next proposition says that, via the Hall inner product, the co-product on  $\Lambda$  (given by plethystic addition of alphabets) induces the natural product on the dual. In fact, this property characterizes the Hall inner product.

**Proposition 1.3.** *If  $\{P_\lambda(X)\}$  and  $\{Q_\lambda(Y)\}$  are a pair of dual bases (with respect to the Hall inner product), and:*

$$P_\lambda(X + X') = \sum_{\mu, \nu} d_{\mu\nu}^\lambda P_\nu(X) P_\mu(X')$$

then:

$$Q_\mu(Y) Q_\nu(Y) = \sum_{\lambda} d_{\mu\nu}^\lambda Q_\lambda(Y)$$

*Proof.* Let:

$$P_\lambda(X + X') = \sum_{\mu, \nu} d_{\mu\nu}^\lambda P_\nu(X) P_\mu(X')$$

and let:

$$Q_\mu(Y) Q_\nu(Y) = \sum_{\lambda} \tilde{d}_{\mu\nu}^\lambda Q_\lambda(Y)$$

We have:

$$\begin{aligned} \Omega((X + X')Y) &= \sum_{\lambda} P_\lambda(X + X') Q_\lambda(Y) \\ &= \sum_{\lambda} Q_\lambda(Y) \sum_{\gamma, \mu} d_{\gamma\mu}^\lambda P_\gamma(X) P_\mu(X') \end{aligned}$$

while:

$$\begin{aligned} \Omega(XY) \Omega(X'Y) &= \left( \sum_{\gamma} P_\gamma(X) Q_\gamma(Y) \right) \left( \sum_{\mu} P_\mu(X') Q_\mu(Y) \right) \\ &= \sum_{\gamma, \mu} P_\gamma(X) P_\mu(X') Q_\gamma(Y) Q_\mu(Y) \\ &= \sum_{\gamma, \mu} P_\gamma(X) P_\mu(X') \sum_{\lambda} \tilde{d}_{\gamma\mu}^\lambda Q_\lambda(Y) \\ &= \sum_{\lambda} Q_\lambda(Y) \sum_{\gamma, \mu} \tilde{d}_{\gamma\mu}^\lambda P_\gamma(X) P_\mu(X') \end{aligned}$$

The result follows now since  $\Omega(XY + X'Y) = \Omega(XY) \Omega(X'Y)$ .  $\square$

**Corollary 1.1** (Taylor’s theorem). *If  $\{P_\lambda(X)\}$  and  $\{Q_\lambda(Y)\}$  are a pair of dual bases then for any symmetric function  $f(X)$  we have:*

$$f(X + X') = \sum_{\gamma} \partial_{Q_\gamma}[f(X)]P_\gamma(X')$$

*Proof.* By linearity, it is sufficient to check on the basis  $\{P_\lambda(X)\}$ , from which the proposition follows by observation that:

$$\begin{aligned} \langle Q_\mu(Y), \partial_{Q_\gamma}[P_\lambda(X)] \rangle &= \langle Q_\mu(Y)Q_\gamma(Y), P_\lambda(X) \rangle \\ &= d_{\mu\gamma}^\lambda \end{aligned}$$

□

Note that as a consequence of Taylor’s theorem, and the self-duality of Schur functions, the fact that:

$$s_\lambda(X + X') = \sum_{\mu} s_{\lambda/\mu}(X)s_\mu(X')$$

implies:

$$s_{\lambda/\mu}(X) = \partial_{s_\mu}[s_\lambda(X)]$$

1.3.2. *Column operators.* Let us introduce now the “translation” and “multiplication” operators of Garsia, Haiman and Tesler [10, 13]:

$$(1) \quad \mathcal{T}(z)[f(X)] = f(X + z)$$

$$(2) \quad \mathcal{P}(z)[f(X)] = \Omega(Xz)f(X)$$

Working with the pair of dual bases  $\{m_\lambda(Y)\}$  and  $\{h_\lambda(X)\}$ , and noting that unless  $\lambda$  is a row partition,  $m_\lambda(X)$  will vanish when  $X$  is an alphabet containing a single letter, we get, by Taylor’s theorem, that:

$$\mathcal{T}(z) = \sum_n z^n \partial_{h_n}$$

Similarly:

$$\mathcal{P}(z) = \sum_n z^n h_n(X)$$

Clearly we have:

$$(3) \quad \mathcal{T}(z) = \mathcal{P}^*(z)$$

Similarly, working with the dual bases  $\{f_\lambda(Y)\}$  and  $\{e_\lambda(X)\}$  and recalling that  $f_\lambda(-Y) = m_\lambda(\epsilon Y)$  we get that:

$$\mathcal{T}(-z) = \sum_n (\epsilon z)^n \partial_{e_n}$$

and:

$$\mathcal{P}(-z) = \sum_n (\epsilon z)^n e_n(X)$$

In other words:

$$\mathcal{P}(-z) = \omega \circ \mathcal{P}(\epsilon z) \circ \omega$$

and:

$$\mathcal{T}(-z) = \omega^* \circ \mathcal{T}(\epsilon z) \circ \omega^*$$

The following lemma tells us how the translation and multiplication operators commute:

**Lemma 1.1.**

$$\mathcal{T}(v)\mathcal{P}(w) = \Omega(vw)\mathcal{P}(w)\mathcal{T}(v)$$

*Proof.*

$$\begin{aligned} \mathcal{T}(v)\mathcal{P}(w)[h(X)] &= \mathcal{T}(v)[\Omega(Xw)h(X)] \\ &= \Omega((X+v)w)h(X+v) \\ &= \Omega(vw)\Omega(Xw)\mathcal{T}(v)[h(X)] \\ &= \Omega(vw)\mathcal{P}(w)\mathcal{T}(v)[h(X)] \end{aligned}$$

□

We can now define the column operator:

$$(4) \quad C(z) = \sum_m (-1)^m C_m z^m = \mathcal{P}(-z)\mathcal{T}(1/z)$$

and the row operator:

$$(5) \quad R(z) = \sum_n z^n (-1)^n R_n = \mathcal{P}(z)\mathcal{T}(-1/z) = C(-z)$$

**Proposition 1.4.** *The Schur function has a natural expression in terms of row operators:*

$$s_\lambda(X) = R_{\lambda_1} \circ R_{\lambda_2} \circ \cdots \circ R_{\lambda_n}[1]$$

*or in terms of column operators:*

$$s_\lambda(X) = C_{\lambda'_1} \circ C_{\lambda'_2} \circ \cdots \circ C_{\lambda'_m}[1]$$

*Proof.* Starting from the definition of the Schur function as:

$$s_\lambda(X) = \det(e_{\lambda'_i + j - i}(X))$$

and expanding along the first row we get that:

$$\begin{aligned} s_\lambda(X) &= \sum_{k=0}^{n-1} (-1)^k e_{\lambda'_1+k}(X) s_{\mu/(k)}(X) \\ &= \sum_{k=0}^{n-1} (-1)^k e_{\lambda'_1+k}(X) \partial_{h_k(X)}[s_\mu(X)] \\ &= C_{\lambda'_1}[s_\mu(X)] \end{aligned}$$

where  $\mu$  is the partition obtained from  $\lambda$  by removing the first column.

Similarly, from the definition of the Schur function as:

$$s_\lambda(X) = \det(h_{\lambda_i+j-i}(X))$$

we get by expanding along the first row that:

$$s_\lambda(X) = R_{\lambda_1}[s_\mu(X)]$$

where this time  $\mu$  is the partition obtained from  $\lambda$  by removing the first row.

□

**Corollary 1.2.** *In the row and column cases we have:*

$$\begin{aligned} s_{(n)}(X) &= h_n(X) \\ s_{(1^n)}(X) &= e_n(X) \end{aligned}$$

1.3.3. *Duality.* The Schur functions are, up to scaling, the unique homogeneous basis which is self dual with respect to the Hall inner product. We give here a slightly non-standard proof of this fact, which will be generalized in the next section.

**Proposition 1.5.** *The Schur functions are self-dual with respect to the Hall inner product*

*Proof.* By proposition 1.3 and corollary 1.2 it suffices to demonstrate that:

$$\langle \Omega_z(Y) s_\mu(Y), s_\lambda(X) \rangle = \langle s_\mu(Y), s_\lambda(X+z) \rangle$$

Begin by observing that:

$$\begin{aligned} \mathcal{T}(v)C(z) &= \mathcal{T}(v)\mathcal{P}(-z)\mathcal{T}(1/z) \\ &= \Omega(-vz)\mathcal{P}(-z)\mathcal{T}(v)\mathcal{T}(1/z) \\ &= (1-vz)C(z)\mathcal{T}(v) \end{aligned}$$

which tells us that:

$$[C(z), \mathcal{T}(v)] = vz C(z) \mathcal{T}(v)$$

Equating coefficients of  $(-1)^m v^k z^m$  on both sides we get that:

$$[C_m, \partial_{h_k}] = -C_{m-1} \partial_{h_{k-1}}$$

Rewriting this in the form:

$$\partial_{h_m} C_k = C_k \partial_{h_m} + C_{k-1} \partial_{h_{m-1}}$$

we obtain:

$$\begin{aligned} \partial_{h_k} [s_\lambda(X)] &= \partial_{h_k} \circ C_{\lambda'_1} \circ C_{\lambda'_2} \circ \cdots \circ C_{\lambda'_n} [1] \\ &= \sum_{\substack{I \subseteq [n] \\ |I|=k}} C_{\lambda'_1 - a_1} \circ C_{\lambda'_2 - a_2} \circ \cdots \circ C_{\lambda'_n - a_n} [1] \end{aligned}$$

where  $a_i = 1$  if  $i \in I$  and zero otherwise.

Of course it is possible that for some  $k$  we have  $\lambda_k = \lambda_{k+1}$  whilst  $a_k \in I$  and  $a_{k-1} \notin I$ , in which case the sequence of integers  $(\lambda_1 - a_1, \lambda_2 - a_2, \dots, \lambda_n - a_n)$  will not form a partition. In this case, however, we have that the determinant:  $\det(e_{\lambda'_i - a_i + j - i}(X))$  contains a repeated column, and thus vanishes.

We conclude that:

$$(6) \quad \mathcal{T}(z)[s_\lambda(X)] = s_\lambda(X + z) = \sum_{\mu \in D(\lambda)} s_\mu(X) z^{|\lambda| - |\mu|}$$

where  $D(\lambda)$  denotes all the partitions which can be obtained from  $\lambda$  by removing a horizontal strip. This is the first form of the recurrence for the Schur functions.

Dually:

$$\begin{aligned} \mathcal{P}(v)C(z) &= \mathcal{P}(v)\mathcal{P}(-z)\mathcal{T}(1/z) \\ &= \mathcal{P}(-z)\mathcal{P}(v)\mathcal{T}(1/z) \\ &= \mathcal{P}(-z)\Omega(-v/z)\mathcal{T}(1/z)\mathcal{P}(-v) \\ &= (1 - v/z)C(z)\mathcal{P}(v) \end{aligned}$$

which tells us that:

$$[C(z), \mathcal{P}(v)] = v/z C(z) \mathcal{P}(v)$$

By equating coefficients of  $(-1)^m v^k z^m$  on both sides we find that:

$$[C_m, h_k(X)] = -C_{m+1} h_{k-1}(X)$$

Rewriting this in the form:

$$h_m(X)C_k = C_k h_m(X) + C_{k+1} h_{m-1}$$

a similar argument to the previous reveals:

$$(7) \quad \mathcal{P}(z)[s_\mu(Y)] = s_\mu(Y)\Omega_z(Y) = \sum_{\lambda \in U(\mu)} s_\lambda(Y)z^{|\lambda|-|\mu|}$$

where  $U(\mu)$  denotes the set of all partitions which can be obtained from  $\mu$  by *adding* a horizontal strip. This is the first form of the Pieri formula for the Schur functions.

Combining these two facts, we see that:

$$\langle s_\mu(Y)\Omega(Yz), s_\lambda(X) \rangle = \langle s_\mu(Y), s_\lambda(X+z) \rangle$$

as claimed. □

Note that identical arguments using the row operator rather than the column operator can be used to show that:

$$(8) \quad \mathcal{T}(-z)[s_\lambda(X)] = s_\lambda(X-z) = \sum_{\mu \in \tilde{D}(\lambda)} s_\mu(X)(\epsilon z)^{|\lambda|-|\mu|}$$

where  $\tilde{D}(\lambda)$  denotes all the partitions which may be obtained from  $\lambda$  by removing a vertical strip. This is the second form of the recurrence for the Schur functions.

Also we have:

$$(9) \quad \mathcal{P}(-z)[s_\mu(Y)] = s_\mu(Y)\tilde{\Omega}_z(Y) = \sum_{\lambda \in \tilde{U}(\mu)} s_\lambda(Y)(\epsilon z)^{|\lambda|-|\mu|}$$

where  $\tilde{U}(\mu)$  denotes the set of all partitions which can be obtained from  $\mu$  by adding a vertical strip. This is the second form of the Pieri formula for the Schur functions.

Putting these two facts together we see that we also have:

$$\langle s_\mu(Y)\tilde{\Omega}(Yz), s_\lambda(X) \rangle = \langle s_\mu(Y), s_\lambda(X-z) \rangle$$



**1.4. Littlewood–Richardson Bases.** The Littlewood–Richardson coefficients  $c_{\mu\nu}^\lambda$  are defined by:

$$s_\lambda(X + X') = \sum_{\mu, \nu} c_{\mu\nu}^\lambda s_\mu(X) s_\nu(X')$$

These numbers may be described nicely by a combinatorial object known as *puzzles* [34]. A very nice proof of the Littlewood–Richardson rule using ideas from Quantum Integrability can be found in [44]. In this section, we describe all the other bases  $\{P_\lambda(X)\}$  of the ring of symmetric functions  $\Lambda$  with the property:

$$P_\lambda(X + X') = \sum_{\mu, \nu} c_{\mu\nu}^\lambda P_\mu(X) P_\nu(X')$$

We shall say that such a basis is of *Littlewood–Richardson type*.

Note that we are only interested in bases for  $\Lambda$  (the ring of symmetric functions in infinitely many variables) which are stable in the sense that if  $\lambda$  is a partition with at most  $k$  parts, its expansion in terms of Schur functions is unchanged under the projection onto  $\Lambda_k$  (the ring of symmetric functions in just  $k$  variables).

We saw in the proof of proposition 1.5 that:

$$s_\lambda(X + z) = \sum_{\mu \in D(\lambda)} s_\mu(X) s_{(|\lambda| - |\mu|)}(z)$$

To prove that a given basis  $\{P_\lambda(X)\}$  is of Littlewood–Richardson type it suffices to demonstrate that:

$$P_\lambda(X + z) = \sum_{\mu \in D(\lambda)} P_\mu(X) P_{(|\lambda| - |\mu|)}(z)$$

Since the Schur functions are self-dual we have, by proposition 1.3, that:

$$s_\mu(Y) s_\nu(Y) = \sum_{\lambda} c_{\mu\nu}^\lambda s_\lambda(Y)$$

If  $\{P_\lambda(X)\}$  is a basis of Littlewood–Richardson type, and  $\{Q_\lambda(Y)\}$  is its dual basis, then we must have:

$$Q_\mu(Y) Q_\nu(Y) = \sum_{\lambda} c_{\mu\nu}^\lambda Q_\lambda(Y)$$

We also saw in the proof of proposition 1.3 that:

$$s_\mu(Y) \sum_n s_{(n)}(Y) s_{(n)}(z) = \sum_{\lambda \in U(\mu)} s_\lambda(Y) s_{(|\lambda| - |\mu|)}(z)$$

This will generalize to:

$$Q_\mu(Y) \sum_n Q_{(n)}(Y) Q_{(n)}(z) = \sum_{\lambda \in U(\mu)} Q_\lambda(Y) Q_{(|\lambda| - |\mu|)}(z)$$

1.4.1. *Generalized complete symmetric functions.* In the row case, the Schur functions reduce to the complete symmetric functions, which have generating function:

$$\Omega(Xz) = \sum_n h_n(X) z^n$$

The complete symmetric functions have the property that:

$$h_n(X + X') = \sum_k h_k(X) h_{n-k}(X')$$

Let  $f(z)$  be an arbitrary delta series, with compositional inverse  $g(z)$ .

Define generalized complete symmetric functions by:

$$(10) \quad \Omega_f(Xz) = \prod_x \frac{1}{1 - f(z)x} = \sum_n r_n(X) z^n$$

Also define, in the dual space:

$$(11) \quad \Phi_g(Yz) = \prod_y \frac{1}{1 - zg(y)} = \sum_n \rho_n(Y) z^n$$

Since:

$$\Omega_f(X + X') = \Omega_f(X) \Omega_f(X')$$

we have that:

$$r_n(X + X') = \sum_k r_k(X) r_{n-k}(X')$$

Furthermore, we have that:

$$r_n(X) = \sum_k \alpha_{nk} h_k(X)$$

where:

$$\alpha_{nk} = [y^n] f(y)^k$$

In other words,  $r_n(X)$  is the image of the convolution type sequence associated to  $f(y)$  under co-algebra embedding of  $\mathbb{Q}[x]$  into  $\Lambda$  given by:

$$\frac{x^n}{n!} \mapsto h_n(X)$$

Note that there are plenty of other collections of ring generators  $\{p_n(X)\}$  for  $\Lambda$  with the property that:

$$p_n(X + X') = \sum_k p_k(X) p_{n-k}(X')$$

For any pair of sequences  $\{f_n(x)\}$  and  $\{g_n(y)\}$  such:

$$\sum_n f_n(x) g_n(y) = \frac{1}{1 - xy}$$

the ring generators  $\{p_n(X)\}$  defined by:

$$\sum_n p_n(X) z^n = \prod_x f_n(x) z^n = \prod_x x^n g_n(z)$$

will have this property. However, unless they are of the special form above, it will not be possible to write them as:

$$p_n(X) = \sum_k \square_{nk} h_k(X)$$

instead they must have more general expansions of the form:

$$p_n(X) = \sum_\lambda \square_\lambda h_\lambda(X)$$

Their image under the projection:

$$h_n(X) \mapsto \frac{x^n}{n!}$$

will still be a sequence of convolution type associated to some  $f(y)$ , however.

The idea now is to show that the same determinantal construction which produces the Schur functions from the complete symmetric functions can be used to produce more general bases with the Littlewood–Richardson property.

In the special case where the ring bases are images of convolution type sequences, we shall see that, as a direct consequence of the multilinearity of the determinant, the associated Littlewood–Richardson bases have expansions of the form:

$$P_\lambda(X) = \sum_{\nu \subseteq \lambda} \square_{\lambda\nu} s_\nu(X)$$

while the dual basis will have an expansion of the form:

$$Q_\mu(Y) = \sum_{\mu \subseteq \nu} \square_{\nu\mu} s_\nu(Y)$$

1.4.2. *Umbral operators.* Let  $U_f$  be the operator which is defined on the complete symmetric functions by:

$$U_f[\Omega(Xz)] = \Omega_f(Xz)$$

extended multiplicatively to the whole of  $\Lambda$ . Clearly we have:

$$U_f^{-1} = U_g$$

It turns out that if  $P_\lambda(X) = U_f[s_\lambda(X)]$  then  $\{P_\lambda(X)\}$  is a basis of Littlewood–Richardson type with dual basis  $\{Q_\lambda(Y)\}$  where  $Q_\lambda(Y) = U_g^*[s_\lambda(Y)]$ .

**Proposition 1.6.**

$$U_g^*[\Omega(Yz)] = \Phi_g(Yz)$$

*Proof.* Let  $\{n_\lambda(Y)\}$  denote the dual basis to  $\{r_\lambda(X)\}$ . We have:

$$\begin{aligned} \langle U_f^*[n_\mu(Y)], h_\lambda(X) \rangle &= \langle n_\mu(Y), U_f[h_\lambda(X)] \rangle \\ &= \langle n_\mu(Y), r_\lambda(X) \rangle \\ &= \delta_{\lambda\mu} \end{aligned}$$

and so:

$$U_f^*[n_\mu(Y)] = m_\mu(Y)$$

That is:

$$n_\mu(Y) = U_g^*[m_\mu(Y)]$$

Now:

$$\begin{aligned} \Omega(XY) &= \prod_y \Omega(Xy) \\ &= \prod_y \sum_n r_n(X) g(y)^n \end{aligned}$$

which gives us the following explicit expression for  $n_\lambda(Y)$ :

$$n_\lambda(Y) = \sum_\sigma \prod_k g(y_k)^{\lambda_{\sigma(k)}}$$

where the sum is over all distinct permutations of  $\lambda$ . Finally:

$$\begin{aligned} U_g^*[\Omega(Yz)] &= \sum_n U_g^*[m_\lambda(Y)] z^{|\lambda|} \\ &= \sum_n n_\lambda(Y) z^{|\lambda|} \\ &= \prod_x \frac{1}{1 - z g(y)} \end{aligned}$$

□

**Proposition 1.7.**      •  $\partial_{r_n} = U_g^* \circ \partial_{h_n} \circ U_f^*$   
 •  $\partial_{\rho_n} = U_g \circ \partial_{h_n} \circ U_f$

*Proof.* For the first part:

$$\begin{aligned}
 \langle \partial_{r_n}[n_\mu(Y)], r_\lambda(X) \rangle &= \langle n_\mu(Y), r_\lambda(X) r_n(X) \rangle \\
 &= \delta_{\mu, n+\lambda} \\
 &= \langle m_\mu(Y), h_\lambda(X) h_n(X) \rangle \\
 &= \langle \partial_{h_n}[m_\mu(Y)], h_\lambda(X) \rangle \\
 &= \langle \partial_{h_n} \circ U_f^*[n_\mu(X)], U_g[r_\lambda(X)] \rangle \\
 &= \langle U_g^* \circ \partial_{h_n} \circ U_f^*[n_\mu(Y)], r_\lambda(X) \rangle
 \end{aligned}$$

For the second part, let  $\{\eta_\lambda(X)\}$  denote the dual basis to  $\{\rho_\lambda(Y)\}$ . As before we have:

$$\begin{aligned}
 \langle h_\mu(Y), U_f[\eta_\lambda(X)] \rangle &= \langle U_f^*[h_\mu(Y)], \eta_\lambda(X), \rangle \\
 &= \langle \rho_\mu(Y), \eta_\lambda(X) \rangle \\
 &= \delta_{\lambda\mu}
 \end{aligned}$$

and so:

$$U_f[\eta_\lambda(X)] = m_\lambda(X)$$

That is:

$$\eta_\lambda(X) = U_g[m_\lambda(X)]$$

Now:

$$\begin{aligned}
 \langle \rho_\mu(Y), \partial_{\rho_n}[\eta_\lambda(X)] \rangle &= \langle \rho_\mu(Y) \rho_{(n)}(Y), \eta_\lambda(X) \rangle \\
 &= \delta_{\mu+(n), \lambda} \\
 &= \langle h_\mu(Y) h_{(n)}(Y), m_\lambda(X) \rangle \\
 &= \langle h_\mu(Y), \partial_{h_n}[m_\lambda(X)] \rangle \\
 &= \langle U_g^*[\rho_\mu(Y)], \partial_{h_n} \circ U_f[\eta_\lambda(X)] \rangle \\
 &= \langle \rho_\mu(Y), U_g \circ \partial_{h_n} \circ U_f[\eta_\lambda(X)] \rangle
 \end{aligned}$$

□

1.4.3. *Column operators.* The (generalized) column operator for the Littlewood–Richardson basis  $\{P_\lambda(X)\}$  is:

$$C_f(z) = \sum_m (-1)^m C_m^f z^m = U_f \circ C(z) \circ U_g$$

where  $C(z)$  is the column operator for the Schur functions.

The generalized column operator for the dual basis  $\{Q_\lambda(Y)\}$  is:

$$\hat{C}_f(z) = \sum_m (-1)^m \hat{C}_m^f z^m = C_f^*(z) = U_g^* \circ C^*(z) \circ U_f^*$$

Since these operators are adjoint, its clear that the resulting bases:

$$P_\lambda(X) = C_{\lambda'_1}^f \circ C_{\lambda'_2}^f \circ \cdots \circ C_{\lambda'_n}^f [1]$$

$$Q_\lambda(Y) = \hat{C}_{\lambda'_1}^f \circ \hat{C}_{\lambda'_2}^f \circ \cdots \circ \hat{C}_{\lambda'_n}^f [1]$$

are dual. More explicitly:

$$\begin{aligned} \langle Q_\mu(Y), P_\lambda(X) \rangle &= \langle U_g^*[s_\mu(Y)], U_f[s_\lambda(X)] \rangle \\ &= \langle s_\mu(Y), U_g \circ U_f[s_\lambda(X)] \rangle \\ &= \langle s_\mu(Y), s_\lambda(X) \rangle \\ &= \delta_{\lambda\mu} \end{aligned}$$

Note that by the same argument used in proposition 1.4 we have:

$$P_\lambda(X) = \det(r_{\lambda_i+j-i}(X))$$

$$Q_\lambda(Y) = \det(\rho_{\lambda_i+j-i}(Y))$$

Now define the generalized multiplication operator:

$$\begin{aligned} \mathcal{P}_f(z) &= \sum_k z^k r_k(X) \\ &= \sum_k f(z)^k h_k(X) \\ &= U_f \circ \mathcal{P}(z) \circ U_g \end{aligned}$$

as well as the generalized translation operator:

$$\begin{aligned} \mathcal{T}_g(z) &= \sum_k z^k \partial_{\rho_k} \\ &= \sum_k r_k(z) \partial_{h_k} \\ &= U_f \circ \mathcal{T}(z) \circ U_g \end{aligned}$$

Also let us define, in the dual space:

$$\begin{aligned}\hat{\mathcal{P}}_g(z) &= \mathcal{T}_g^*(z) \\ &= U_g^* \circ \mathcal{P}(z) \circ U_f^* \\ &= \sum_k z^k \rho_k(Y) \\ &= \sum_k r_k(z) h_k(X)\end{aligned}$$

as well as:

$$\begin{aligned}\hat{\mathcal{T}}_f(z) &= \mathcal{P}_f^*(z) \\ &= U_g^* \circ \mathcal{T}(z) \circ U_f^* \\ &= \sum_k z^k \partial_{r_k} \\ &= \sum_k f(z)^k \partial_{h_k}\end{aligned}$$

We have now that:

$$C_f(z) = \mathcal{P}_{-f}(z) \circ \mathcal{T}_{1/g}(z)$$

while:

$$\hat{C}_f(z) = \hat{\mathcal{P}}_{-f}(z) \circ \hat{\mathcal{T}}_{1/g}(z)$$

**Proposition 1.8.** *The (non-homogeneous) basis for the ring of symmetric functions defined by:*

$$P_\lambda(X) = C_{\lambda'_1}^f \circ C_{\lambda'_2}^f \circ \cdots \circ C_{\lambda'_m}^f [1]$$

*is of Littlewood–Richardson type.*

*Proof.* We must show that:

$$\langle Q_\lambda(Y) \Phi_g(Yz), P_\mu(X) \rangle = \langle Q_\lambda(Y), P_\mu(X + z) \rangle$$

By conjugating Lemma 1.1 by  $U_f$  on the left and  $U_g$  on the right, we find that:

$$\mathcal{T}_g(v) \circ \mathcal{P}_f(w) = \Omega(vw) \circ \mathcal{P}_f(w) \circ \mathcal{T}_g(v)$$

This in turn implies that:

$$[\tilde{C}(z), \mathcal{T}_g(v)] = vz \circ \tilde{C}(z) \circ \mathcal{T}_g(v)$$

By arguments which are essentially identical to those in the proof of proposition 1.5 we obtain the following generalized recurrence:

$$\partial_{\rho_k}[P_\lambda(X)] = \sum_{\mu \in D_k(\lambda)} P_\mu(X)$$

Next by using the fact that:

$$\mathcal{T}(z) = \sum_k r_k(z) \partial_{\rho_k}$$

we can re-write the above as:

$$\mathcal{T}(z)[P_\lambda(X)] = \sum_{\mu \in D(\lambda)} P_\mu(X) P_{(|\lambda| - |\mu|)}(z)$$

Similarly, by conjugating Lemma 1.1 on the left by  $U_g^*$  and the right by  $U_f^*$  we find that:

$$\hat{\mathcal{T}}_g(v) \circ \hat{\mathcal{P}}_f(w) = \Omega(vw) \circ \hat{\mathcal{P}}_f(w) \circ \hat{\mathcal{T}}_g(v)$$

This tells us that:

$$[\hat{C}(z), \hat{\mathcal{P}}_f(v)] = v/z \circ \hat{C}(z) \circ \hat{\mathcal{P}}_f(v)$$

and so we have the following Pieri formula :

$$Q_\mu(Y) \rho_k(Y) = \sum_{\lambda \in U_k(\mu)} Q_\lambda(Y)$$

That is:

$$\hat{\mathcal{P}}_f(z)[Q_\mu(Y)] = Q_\mu(Y) \Phi_g(Yz) = \sum_{\lambda \in U(\mu)} Q_\lambda(Y) z^{|\lambda| - |\mu|}$$

Putting these two facts together we find that:

$$\langle Q_\lambda(Y) \Phi_g(Yz), P_\mu(X) \rangle = \langle Q_\lambda(Y), P_\mu(X + z) \rangle$$

as claimed.  $\square$

1.4.4. *Generalized elementary symmetric functions.* Now define generalized elementary symmetric functions:

$$\tilde{\Omega}_f(Xz) = \prod_x (1 - f(-z)x) = \sum_n c_n(X) z^n$$

and in the dual space:

$$\tilde{\Phi}_g(Yz) = \prod_y (1 + zg(y)) = \sum_n \gamma_n(Y) z^n$$

A minor modification of the argument at the end of section 1.1.4 can be used to show that:

$$\det(r_{\lambda_i + j - i}(X)) = \det(c_{\lambda'_i + j - i}(\epsilon X))$$

$$\det(\rho_{\lambda_i + j - i}(Y)) = \det(\gamma_{\lambda'_i + j - i}(\epsilon Y))$$



The generalized row operators are:

$$R_f(z) = C_f(-z)$$

$$\hat{R}_f(z) = \hat{C}_f(-z)$$

One can show that:

$$P_\lambda(X) = R_{\lambda_1}^f \circ R_{\lambda_2}^f \circ \cdots \circ R_{\lambda_n}^f[1]$$

$$Q_\lambda(Y) = \hat{R}_{\lambda_1}^f \circ \hat{R}_{\lambda_2}^f \circ \cdots \circ \hat{R}_{\lambda_n}^f[1]$$

which implies that:

$$\mathcal{T}(-z)[P_\lambda(X)] = P_\lambda(X - z) = \sum_{\mu \in \tilde{D}_\lambda} P_\mu(X) P_{(|\lambda| - |\mu|)}(\epsilon z)$$

$$\hat{\mathcal{P}}_{-f}(z)[Q_\mu(Y)] = Q_\mu(Y) \tilde{\Phi}_f(Yz) = \sum_{\lambda \in \tilde{U}(\mu)} Q_\lambda(Y) Q_{(|\lambda| - |\mu|)}(\epsilon z)$$

In other words:

$$\langle Q_\lambda(Y) \tilde{\Phi}_f(Yz), P_\mu(X) \rangle = \langle Q_\lambda(Y), P_\mu(X - z) \rangle$$

**1.5. Examples.** In this section we shall give symmetric function analogues of the rising and falling factorials, as well as the rook polynomials, leading to a generalized family of Stirling and Lah numbers which are indexed by partitions rather than integers.

For any formal power series  $f(z)$  let  $\bar{f}(z) = f(-z)$ .

Let us use the following notation for the falling factorial:

$$(x)_n = x(x-1)(x-2) \cdots (x-n+1)$$

and the rising factorial:

$$(x)^n = x(x+1)(x+2) \cdots (x+n-1)$$

The Stirling numbers of the first kind are defined by:

$$(x)_n = \sum_k s(n, k) x^k$$

We have:

$$(x)^n = \sum_k |s(n, k)| x^k$$

One may check that  $\{\frac{(x)_n}{n!}\}$  is the sequence of convolution type associated to  $f(z) = (\exp(z) - 1)$  while  $\{\frac{(x)^n}{n!}\}$  is the sequence of convolution type associated to  $-\bar{f}(z) = (1 - \exp(-z))$ .

Now let  $\{A_\lambda(X)\}$  denote the Littlewood–Richardson basis for the ring of symmetric functions  $\Lambda$  that is associated to  $f(z) = \exp(z) - 1$ . The generating function for the row case is:

$$\Omega_f(Xz) = \prod_{x \in X} \frac{1}{1 - x \log(1 + z)}$$

while the generating function for the column case is:

$$\tilde{\Omega}_f(Xz) = \prod_{x \in X} (1 - x \log(1 - z))$$

Similarly let  $\{B_\lambda(X)\}$  denote the Littlewood–Richardson basis associated to  $-\bar{f}(z) = 1 - \exp(-z)$ . The generating function for the row case is:

$$\begin{aligned} \Omega_{-\bar{f}}(Xz) &= \prod_{x \in X} \frac{1}{1 - x \log\left(\frac{1}{1-z}\right)} \\ &= \prod_{x \in X} \frac{1}{1 + x \log(1 - z)} \\ &= \tilde{\Omega}_f(-\epsilon Xz) \\ &= \omega[\tilde{\Omega}_f(Xz)] \end{aligned}$$

The generating function for the column case is:

$$\begin{aligned} \tilde{\Omega}_{-\bar{f}}(Xz) &= \prod_{x \in X} \left(1 - x \log\left(\frac{1}{1+z}\right)\right) \\ &= \prod_{x \in X} (1 + x \log(1 + z)) \\ &= \Omega_f(-\epsilon Xz) \\ &= \omega[\Omega_f(Xz)] \end{aligned}$$

Let  $A = (a_{\lambda\mu})$  denote the transition matrix from the basis  $\{A_\lambda(X)\}$  to the usual Schur basis  $\{s_\lambda(X)\}$ , and let  $B = (b_{\lambda\mu})$  denote the transition matrix from  $\{B_\lambda(X)\}$  to  $\{s_\lambda(X)\}$ . On the next page we give the corner of these matrices corresponding to partitions with at most five parts.

The order which we are using on partitions is:

$$\begin{aligned}
& \{\{1\}\} \\
& \{\{2\}, \{1, 1\}\} \\
& \{\{3\}, \{2, 1\}, \{1, 1, 1\}\} \\
& \{\{4\}, \{3, 1\}, \{2, 2\}, \{2, 1, 1\}, \{1, 1, 1, 1\}\} \\
& \{\{5\}, \{4, 1\}, \{3, 2\}, \{3, 1, 1\}, \{2, 2, 1\}, \{2, 1, 1, 1\}, \{1, 1, 1, 1, 1\}\}
\end{aligned}$$

Here is a portion of the matrix  $A$ :

$$\begin{pmatrix}
1 & -\frac{1}{2} & \frac{1}{2} & \frac{1}{3} & -\frac{1}{3} & \frac{1}{3} & -\frac{1}{4} & \frac{1}{4} & 0 & -\frac{1}{4} & \frac{1}{4} & \frac{1}{5} & -\frac{1}{5} & 0 & \frac{1}{5} & 0 & -\frac{1}{5} & \frac{1}{5} \\
0 & 1 & 0 & -1 & \frac{1}{2} & 0 & \frac{11}{12} & -\frac{7}{12} & -\frac{1}{12} & \frac{1}{3} & 0 & -\frac{5}{6} & \frac{7}{12} & \frac{1}{12} & -\frac{5}{12} & -\frac{1}{12} & \frac{1}{4} & 0 \\
\vdots & \ddots & 1 & 0 & -\frac{1}{2} & 1 & 0 & \frac{1}{3} & -\frac{1}{12} & -\frac{7}{12} & \frac{11}{12} & 0 & -\frac{1}{4} & \frac{1}{12} & \frac{5}{12} & -\frac{1}{12} & -\frac{7}{12} & \frac{5}{6} \\
& & & 1 & 0 & 0 & -\frac{3}{2} & \frac{1}{2} & 0 & 0 & 0 & \frac{7}{4} & -\frac{5}{6} & -\frac{1}{12} & \frac{1}{3} & 0 & 0 & 0 \\
& & & & 1 & 0 & 0 & -1 & 0 & 1 & 0 & 0 & \frac{11}{12} & -\frac{1}{12} & -\frac{13}{12} & -\frac{1}{12} & \frac{11}{12} & 0 \\
& & & & & 1 & 0 & 0 & 0 & -\frac{1}{2} & \frac{3}{2} & 0 & 0 & 0 & \frac{1}{3} & -\frac{1}{12} & -\frac{5}{6} & \frac{7}{4} \\
& & & & & & 1 & 0 & 0 & 0 & 0 & -2 & \frac{1}{2} & 0 & 0 & 0 & 0 & 0 \\
& & & & & & & 1 & 0 & 0 & 0 & 0 & -\frac{3}{2} & 0 & 1 & 0 & 0 & 0 \\
& & & & & & & & 1 & 0 & 0 & 0 & 0 & -1 & 0 & 1 & 0 & 0 \\
& & & & & & & & & 1 & 0 & 0 & 0 & 0 & -1 & 0 & \frac{3}{2} & 0 \\
& & & & & & & & & & 1 & 0 & 0 & 0 & 0 & 0 & -\frac{1}{2} & 2 \\
& & & & & & & & & & & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\
& & & & & & & & & & & & 1 & 0 & 0 & 0 & 0 & 0 \\
& & & & & & & & & & & & & 1 & 0 & 0 & 0 & 0 \\
& & & & & & & & & & & & & & 1 & 0 & 0 & 0 \\
& & & & & & & & & & & & & & & \ddots & 1 & 0 \\
& & & & & & & & & & & & & & & \dots & 0 & 1
\end{pmatrix}$$

The columns are the images of the Schur functions under the transformation  $h_k \mapsto \sum_{i \leq k} c_{k,i} h_i$  where  $c_{k,i}$  are the coefficients of the powers of  $\log(1+z)$ .

Here is a portion of the matrix  $B$ :

$$\begin{pmatrix} 1 & \frac{1}{2} & -\frac{1}{2} & \frac{1}{3} & -\frac{1}{3} & \frac{1}{3} & \frac{1}{4} & -\frac{1}{4} & 0 & \frac{1}{4} & -\frac{1}{4} & \frac{1}{5} & -\frac{1}{5} & 0 & \frac{1}{5} & 0 & -\frac{1}{5} & \frac{1}{5} \\ 0 & 1 & 0 & 1 & -\frac{1}{2} & 0 & \frac{11}{12} & -\frac{7}{12} & -\frac{1}{12} & \frac{1}{3} & 0 & \frac{5}{6} & -\frac{7}{12} & -\frac{1}{12} & \frac{5}{12} & \frac{1}{12} & -\frac{1}{4} & 0 \\ \vdots & \ddots & 1 & 0 & \frac{1}{2} & -1 & 0 & \frac{1}{3} & -\frac{1}{12} & -\frac{7}{12} & \frac{11}{12} & 0 & \frac{1}{4} & -\frac{1}{12} & -\frac{5}{12} & \frac{1}{12} & \frac{7}{12} & -\frac{5}{6} \\ \hline & & & 1 & 0 & 0 & \frac{3}{2} & -\frac{1}{2} & 0 & 0 & 0 & \frac{7}{4} & -\frac{5}{6} & -\frac{1}{12} & \frac{1}{3} & 0 & 0 & 0 \\ & & & & 1 & 0 & 0 & 1 & 0 & -1 & 0 & 0 & \frac{11}{12} & -\frac{1}{12} & -\frac{13}{12} & -\frac{1}{12} & \frac{11}{12} & 0 \\ & & & & & 1 & 0 & 0 & 0 & \frac{1}{2} & -\frac{3}{2} & 0 & 0 & 0 & \frac{1}{3} & -\frac{1}{12} & -\frac{5}{6} & \frac{7}{4} \\ \hline & & & & & & 1 & 0 & 0 & 0 & 0 & 2 & -\frac{1}{2} & 0 & 0 & 0 & 0 & 0 \\ & & & & & & & 1 & 0 & 0 & 0 & 0 & \frac{3}{2} & 0 & -1 & 0 & 0 & 0 \\ & & & & & & & & 1 & 0 & 0 & 0 & 0 & 1 & 0 & -1 & 0 & 0 \\ & & & & & & & & & 1 & 0 & 0 & 0 & 0 & 1 & 0 & -\frac{3}{2} & 0 \\ & & & & & & & & & & 1 & 0 & 0 & 0 & 0 & 0 & \frac{1}{2} & -2 \\ \hline & & & & & & & & & & & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ & & & & & & & & & & & & 1 & 0 & 0 & 0 & 0 & 0 \\ & & & & & & & & & & & & & 1 & 0 & 0 & 0 & 0 \\ & & & & & & & & & & & & & & 1 & 0 & 0 & 0 \\ & & & & & & & & & & & & & & & \ddots & 1 & 0 \\ \vdots & & & & & & & & & & & & & & & & \ddots & 1 & 0 \\ 0 & \dots & & & & & & & & & & & & & & & \dots & 0 & 1 \end{pmatrix}$$

A similar remark applies to  $B$  with  $\log(1+z)$  replaced with  $-\log(1-z)$ . Observe that  $b_{\lambda\mu} = a_{\lambda'\mu'}$ , and  $b_{\lambda\mu} = (-1)^{|\lambda|-|\mu|}a_{\lambda\mu}$ .

If one looks at just entries of  $A$  corresponding to row diagrams (scaled by  $n!/k!$ ), or alternatively those of  $B$  corresponding to column diagrams, one recovers the Stirling numbers of the first kind:

$$\begin{pmatrix} 1 & -1 & 2 & -6 & 24 \\ 0 & 1 & -3 & 11 & -50 \\ 0 & 0 & 1 & -6 & 35 \\ 0 & 0 & 0 & 1 & -10 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

Switching the roles of  $A$  and  $B$ , one obtains the unsigned Stirling numbers of the first kind:

$$\begin{pmatrix} 1 & 1 & 2 & 6 & 24 \\ 0 & 1 & 3 & 11 & 50 \\ 0 & 0 & 1 & 6 & 35 \\ 0 & 0 & 0 & 1 & 10 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

Let us define the Lah numbers by:

$$(x)_n = \sum_k L(n, k)(x)^k$$

Let  $h(z) = \frac{z}{1+z}$  and observe that  $h^{-1}(z) = \frac{z}{1-z}$   
We have:

$$\begin{aligned} -\bar{g}(h(z)) &= \log \left( \frac{1}{1 - \frac{z}{z+1}} \right) \\ &= \log(1 + z) \\ &= g(z) \end{aligned}$$

In other words:

$$U_{-\bar{f}} \circ U_{h^{-1}} = U_f$$

Suppose now that the rook polynomials have the expansion:

$$l_n(x) = U_{h^{-1}} \left[ \frac{x^n}{n!} \right] = \sum_k a_{nk} \frac{x^k}{k!}$$

then we have:

$$U_{-\bar{f}} [l_n(x)] = \sum_k a_{nk} \frac{(x)^k}{k!}$$

while:

$$U_f \left[ \frac{x^n}{n!} \right] = \frac{(x)_n}{n!}$$

Thus we must have:

$$l_n(x) = \sum_k |L(n, k)| x^k$$

Now consider the Littlewood–Richardson basis  $\{L_\lambda(X)\}$  associated to  $h(z) = \frac{z}{1-z}$ . Here is a portion of the transition matrix  $L = (l_{\lambda\mu})$ :

$$\left( \begin{array}{c|c|c|c|c} 1 & 1 & -1 & 1 & -1 & 1 & -1 & 0 & 1 & -1 & 1 \\ \hline 0 & 1 & 0 & 2 & -1 & 0 & 3 & -2 & 0 & 1 & 0 \\ \hline \vdots & \ddots & 1 & 0 & 1 & -2 & 0 & 1 & 0 & -2 & 3 \\ \hline & & & 1 & 0 & 0 & 3 & -1 & 0 & 0 & 0 \\ & & & & 1 & 0 & 0 & 2 & 0 & -2 & 0 \\ & & & & & 1 & 0 & 0 & 0 & 1 & -3 \\ \hline & & & & & & 1 & 0 & 0 & 0 & 0 \\ & & & & & & & 1 & 0 & 0 & 0 \\ & & & & & & & & 1 & 0 & 0 \\ & & & & & & & & & 1 & 0 \\ & & & & & & & & & & \ddots \\ \hline 0 & \dots & & & & & & & & & \dots \end{array} \right)$$

The columns are the images under the Schur functions under the transformation  $h_k \mapsto \sum_{i \leq k} \binom{k}{i} h_i$ .

Looking at just the rows (scaled by  $n!/k!$ ), we recover the Lah numbers:

$$\begin{pmatrix} 1 & 2 & 6 & 24 & 120 \\ 0 & 1 & 6 & 36 & 240 \\ 0 & 0 & 1 & 12 & 120 \\ 0 & 0 & 0 & 1 & 20 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

One may check that  $AL = B$ .

## 2. A GENERATING FUNCTION IDENTITY FOR MACDONALD POLYNOMIALS

The Macdonald polynomials are a  $qt$ -deformation of the Schur functions. The main result of the second part of this thesis is the proof of a generating function identity for Macdonald polynomials which was originally conjectured by Kawanaka [21].

In section 2.1 we define the Macdonald polynomials and state the Kawanaka conjecture. Section 2.2 contains some technical lemmas which are not needed until the final step of the proof. In section 2.3 we discuss the Pieri formula and recurrence for the Macdonald formulas, which generalize those for the Schur functions which were discussed in detail in the first part of this thesis. The proof itself is contained in section 2.4.

### 2.1. Macdonald Polynomials.

2.1.1. *Notation.* Recall from section 1.1.3 the generating series for the complete symmetric functions:

$$\Omega(zX) = \prod_{x \in X} \frac{1}{1 - zx}$$

Here we again use the plethystic notation. Inside a symmetric function the expression  $\frac{1-t}{1-q}$  denotes the alphabet:

$$\frac{1-t}{1-q} = \{1, q, q^2, \dots - t, -tq, -tq^2, \dots\}$$

Recall also the important distinction between plethystic negation:

$$\Omega(-X) = \prod_{x \in X} (1 - x)$$

and formal negation:

$$\Omega(\epsilon X) = \prod_{x \in X} \frac{1}{1 + x}$$

We remark that the operator  $\Omega$  acting on alphabets plays a role in symmetric function theory that is in many ways analogous to that of the exponential function in the theory of functions of a single variable. In particular:

$$\Omega(X + Y) = \Omega(X)\Omega(Y)$$

Also note that:

$$\Omega\left(\frac{z}{1-q}\right) = \frac{1}{(z; q)_\infty} = \sum_k \frac{z^k}{(q; q)_k}$$

is the first  $q$ -exponential function, mentioned in section 1.1.1, whilst:

$$\Omega\left(\frac{-z}{1-q}\right) = (-z; q)_\infty = \sum_k q^{\binom{k}{2}} \frac{z^k}{(q; q)_k}$$

is the second  $q$ -exponential function.

2.1.2. *Operator definition.* In this section we shall be working with a finite alphabet  $X_n$  with exactly  $n$  letters:

$$X_n = x_1 + x_2 + \cdots + x_n$$

Let  $\Delta(X_n)$  denote the Vandemonde determinant:

$$\Delta(X_n) = \left\| \begin{array}{cccc} 1 & 1 & \cdots & 1 \\ x_1 & x_2 & \cdots & x_n \\ \cdots & \cdots & \cdots & \cdots \\ x_1^{n-1} & x_2^{n-1} & \cdots & x_n^{n-1} \end{array} \right\| = \prod_{i < j} (x_j - x_i)$$

The Macdonald polynomials  $\{P_\lambda(X_n)\}$  were originally introduced in [28] as the eigenfunctions of the operator  $D$ :

$$f(x_1, x_2, \dots, x_n) \mapsto \sum_{i=1}^n \prod_{j \neq i} \frac{(tx_i - x_j)}{(x_i - x_j)} f(x_1, \dots, qx_i, \dots, x_n)$$

with eigenvalues:

$$\sum_{i=1}^n q^{\lambda_i} t^{n-i}$$

We may rewrite this operator using the plethystic notation [24, 13, 10] as:

$$f(X_n) \mapsto \frac{1}{\Delta(X_n)} \sum_{i=1}^n \Delta(X_n + (t-1)x_i) f(X_n + (q-1)x_i)$$

Although it is not possible to take the limit  $n \rightarrow \infty$  without the eigenvalues diverging, by using a slightly modified version of this operator, namely:

$$f(X_n) \mapsto t^{-n} \frac{1}{\Delta(X_n)} \sum_{i=1}^n \left( \Delta(X_n + (t-1)x_i) f(X_n + (q-1)x_i) \right) - \sum_{i=1}^n t^{-i} f(X_n)$$

which has eigenvalues  $\sum_{i=1}^n (q^{\lambda_i} - 1)t^{-i}$ , Macdonald was able to show [29] (page 321) that his polynomials have the property that:

$$P_\lambda(X_n) = \pi_n[P_\lambda(X_{n+1})]$$

where  $\pi_n : \Lambda_{n+1} \rightarrow \Lambda_n$  is the map which sets the  $(n+1)$ th variable equal to zero.



2.1.3. *Characterization using the inner product.* Macdonald showed in [28] that his operator  $D$  is self-adjoint (and thus its eigenfunctions are orthogonal) with respect to the deformed Hall inner product:

$$\langle f(Y), g(X) \rangle_{q,t} = \left\langle f \left( Y \frac{1-t}{1-q} \right), g(X) \right\rangle$$

Dual bases  $\{P_\lambda(X)\}$  and  $\{Q_\lambda(Y)\}$  with respect to the Macdonald inner product are characterized by the property that:

$$\sum_{\lambda} P_{\lambda}(X) Q_{\lambda}(Y) = \Omega \left( XY \frac{1-t}{1-q} \right) = \prod_{\substack{x \in X \\ y \in Y}} \frac{(txy, q)_{\infty}}{(xy, q)_{\infty}}$$

Note the important fact that:

$$\Omega \left( (X + Y) \frac{1-t}{1-q} \right) = \Omega \left( X \frac{1-t}{1-q} \right) \Omega \left( Y \frac{1-t}{1-q} \right)$$

The Macdonald polynomials may be characterized as the unique basis for the ring of symmetric functions over  $\mathbb{Q}(q, t)$  which is both orthogonal (but not orthonormal) with respect to the Macdonald inner product, and whose expansion in terms of the monomial symmetric functions is strictly upper triangular with respect to the dominance order [28] on partitions:

$$P_{\lambda}(X) = \sum_{\mu \triangleleft \lambda} \square_{\lambda\mu} m_{\mu}(X)$$

In particular in the column case the Macdonald polynomials correspond to the elementary symmetric functions:

$$P_{(1^n)}(X) = e_n(X)$$

2.1.4. *Arms and legs.* For  $s \in \lambda$  some box in a partition  $\lambda$  the *arm length*  $a_{\lambda}(s)$  is defined to be the number of boxes in  $\lambda$  lying directly to the right of the box  $s$ , while the *leg length* is defined to be the number of boxes in the partition  $\lambda$  lying directly below the box  $s$ .



In other words, if  $s = (i, j)$  then  $a_{\lambda}(s) = \lambda_i - j$  and  $l_{\lambda}(s) = \lambda'_j - i$ . If the box  $s$  lies outside the partition  $\lambda$  then we defined  $a_{\lambda}(s) = l_{\lambda}(s) = 0$ .

Let  $n(\lambda) = \sum_{s \in \lambda} a_{\lambda}(s)$  and let  $\tilde{n}(\lambda) = \sum_{s \in \lambda} l_{\lambda}(s) = n(\lambda')$ .

Let us define:

$$B_\lambda(q, t) = \sum_{\lambda} q^{a_\lambda(s)} t^{l_\lambda(s)}$$

We have, of course, that:

$$B_{\lambda'}(t, q) = B_\lambda(q, t)$$

It is a surprising fact [29] (pages 338,339) that:

$$\langle P_\lambda(Y), P_\lambda(X) \rangle_{q,t} = \frac{\prod_{s \in \lambda} (1 - q^{a_\lambda(s)+1} t^{l_\lambda(s)})}{\prod_{s \in \lambda} (1 - q^{a_\lambda(s)} t^{l_\lambda(s)+1})} = \Omega((t - q)B_\lambda(q, t))$$

2.1.5. *Duality.* The Macdonald  $Q$ -functions are defined to be dual to the Macdonald  $P$ -functions. Since the Macdonald  $P$ -functions are orthogonal with respect to the Hall inner product we have:

$$Q_\lambda(X) = \frac{P_\lambda(X)}{\langle P_\lambda(Y) P_\lambda(X) \rangle_{q,t}}$$

The operator  $\omega_{q,t}$  given by:

$$\omega_{q,t}[f(X)] = \omega f\left(\frac{1-q}{1-t}X\right)$$

is self-adjoint with respect to the Macdonald inner product.

It is another surprising fact [29] (page 327) that:

$$\omega_{q,t}[P_\lambda(X; q, t)] = Q_{\lambda'}(X; t, q)$$

Using this, one can show that in the one row case, the Macdonald polynomials correspond to the modified version of the complete symmetric functions [29] (page 311) defined by:

$$P_{(n)}(X) = \frac{(q; q)_n}{(t; q)_n} g_n(X)$$

Where:

$$\sum_n g_n(X) z^n = \Omega_z \left( X \frac{1-t}{1-q} \right)$$

2.1.6. *Kawanaka conjecture.* In part II of this thesis we shall prove the following generating function identity for the Macdonald polynomials, which was originally conjectured by Kawanaka [21] and proved in the case of Hall-Littlewood Polynomials ( $q = 0$ ):

$$\sum_{\lambda} \left( \prod_{s \in \lambda} \frac{1 + q^{a_\lambda(s)} t^{l_\lambda(s)+1}}{1 - q^{a_\lambda(s)+1} t^{l_\lambda(s)}} \right) P_\lambda(X; q^2, t^2) = \prod_i \frac{(-tx_i; q)_\infty}{(x_i; q)_\infty} \prod_{i < j} \frac{(t^2 x_i x_j; q^2)_\infty}{(x_i x_j; q^2)_\infty}$$

By  $P_\lambda(X; q^2, t^2)$  we mean the Macdonald polynomial  $P_\lambda(X)$  for which every occurrence of the variable  $q$  has been replaced by  $q^2$  and similarly every occurrence of the variable  $t$  has been replaced by  $t^2$ .

This identity complements the following two generating function identities for Macdonald polynomials which can be found in the case where  $b = 0$  or  $b = 1$  on page 349 of Macdonald [29]:

$$\sum_{\lambda} b^{c(\lambda)} \prod_{\substack{s \in \lambda \\ l_{\lambda}(s) \text{ even}}} \frac{1 - q^{a_{\lambda}(s)} t^{l_{\lambda}(s)+1}}{1 - q^{a_{\lambda}(s)+1} t^{l_{\lambda}(s)}} P_{\lambda}(X; q, t) = \prod_{i \leq j} \prod_i \frac{(bt x_i; q)_{\infty}}{(bx_i)_{\infty}} \frac{(tx_i x_j; q)_{\infty}}{(x_i x_j; q)_{\infty}}$$

$$\sum_{\lambda} b^{r(\lambda)} \prod_{\substack{s \in \lambda \\ l_{\lambda}(s) \text{ even}}} \frac{1 - q^{a_{\lambda}(s)} t^{l_{\lambda}(s)+1}}{1 - q^{a_{\lambda}(s)+1} t^{l_{\lambda}(s)}} P_{\lambda}(X; q, t) = \prod_{i \leq j} \prod_i \frac{(bt x_i; q)_{\infty}}{(bx_i)_{\infty}} \frac{(tx_i x_j; q)_{\infty}}{(x_i x_j; q)_{\infty}}$$

Here  $c(\lambda)$  and  $r(\lambda)$  are the number of columns and rows of odd length, respectively.

More recently some extension of the Hall-Littlewood version of the Kawanaka identity have been proved in [27] but no proof of the more general identity has yet appeared in the literature. The results in the second part of this thesis will eventually be published in [35].

We may rewrite the identity to be proved in the plethystic notation as:

$$\sum_{\lambda} \Omega((q - \epsilon t)) B_{\lambda}(q, t) P_{\lambda}(X; q^2, t^2) = \Omega \left( \frac{1 - \epsilon t}{1 - q} X + \frac{1 - t^2}{1 - q^2} e_2(X) \right)$$

By  $e_2(X)$  we mean the alphabet:

$$e_2(X) = \sum_{i < j} x_i x_j$$

When  $q = \epsilon t$  the identity reduces to the well-known generating function for the Schur functions:

$$\sum_{\lambda} s_{\lambda}(X) = \Omega(X + e_2(X))$$

which is perhaps more familiar in the form:

$$\sum_{\lambda} s_{\lambda}(x_1, \dots, x_n) = \prod_{i=1}^n \frac{1}{1 - x_i} \prod_{1 \leq i < j \leq n} \frac{1}{1 - x_i x_j}$$

The proof makes use of the Pieri rule and the recurrence for Macdonald polynomials, which generalize those for the Schur functions discussed in part one, followed by some combinatorial manipulations, to reduce

the identity to a rational function in the variables  $a_k = q^{\mu_k} t^{m-k}$ . The final step is an induction on the residues at the poles.

**2.2. Resultants.** The *resultant* is defined by:

$$R(Z : A) = \prod_{z \in Z} \prod_{a \in A} (z - a)$$

By the fundamental theorem of algebra, up to a scalar multiple, any polynomial  $p(z)$  may be written in the form:

$$p(z) = R(z : A)$$

where  $A$  is the alphabet of zeros. More generally, any rational function  $r(z) = \frac{p(z)}{q(z)}$  may be written, up to a scalar multiple, in the form:

$$r(z) = \frac{R(z : A)}{R(z : B)}$$

where  $A$  is the alphabet of zeros  $B$  is the set of poles.

We shall need to make use of an extended version of the resultant defined by:

$$R(X - Y : A - B) = \frac{R(X : A - B)}{R(Y : A - B)} = \frac{R(X : A)R(Y : B)}{R(Y : A)R(X : B)}$$

Be warned that:

$$\Omega(X - Y) = R(1 : Y - X)$$

In the previous lemma we assumed that the alphabets  $A$  and  $B$  were distinct. In what follows we shall need to consider the case where one alphabet is a scalar multiple of the other.

Let us define:

$$W(X : Y)_{q,t} = R(X : (q/t - 1)Y) = \prod_{\substack{x \in X \\ y \in Y}} \frac{(x - qy/t)}{(x - y)}$$

$$V(X : Y)_{q,t} = R(X : (t/q - 1)Y) = \prod_{\substack{x \in X \\ y \in Y}} \frac{(x - ty/q)}{(x - y)}$$

Observe that:

$$V(X : Y)_{q,t} = W(X : Y)_{1/q, 1/t} = W(X : Y)_{t,q}$$

We shall also need to make use of the functions:

$$v(X : Y)_{q,t} = V(X : qY)_{q,t} = R(X : (t - q)Y) = \prod_{\substack{x \in X \\ y \in Y}} \frac{(x - ty)}{(x - qy)}$$

$$w(X : Y)_{q,t} = W(X : Y/q)_{q,t} = R(X : (1/t - 1/q)Y) = \prod_{\substack{x \in X \\ y \in Y}} \frac{(x - y/t)}{(x - y/q)}$$

Observe that:

$$v(X : Y)_{q,t}^{-1} = W(X : tY)$$

$$w(X : Y)_{q,t}^{-1} = V(X : Y/t)$$

Now let us define:

$$\begin{aligned} \Theta(X : Y)_{q,t} &= v(X : Y)_{q,t} W(X : Y)_{q,t} \\ &= R(X : ((t - q) + (q/t - 1))Y) \\ &= \prod_{\substack{x \in X \\ y \in Y}} \frac{(x - ty)(x - qy/t)}{(x - qy)(x - y)} \end{aligned}$$

as well as:

$$\begin{aligned} \Phi(X : Y)_{q,t} &= V(X : Y)_{q,t} w(X : Y)_{q,t} \\ &= R(X : ((t/q - 1) + (1/t - 1/q))Y) \\ &= \prod_{\substack{x \in X \\ y \in Y}} \frac{(x - ty/q)(x - y/t)}{(x - y)(x - y/q)} \end{aligned}$$

Observe that:

$$\Phi(X : Y)_{q,t} = \Theta(Y : X)_{q,t} = \Theta(X : Y)_{1/q, 1/t} = \Theta(qX : Y)_{q,t}$$

These rational functions will play a key role in the proof of the Kawanaka conjecture. We shall suppress the subscripts when no ambiguity can result.

### 2.2.1. Residue calculations.

**Lemma 2.1.** *For any  $k \geq 1$  we have:*

$$\sum_{\substack{X' + X'' = X \\ |X'| = k}} (\Phi(X' : X'') - \Phi(X'' : X')) = 0$$

*Proof.* We wish to show that for all  $1 \leq k < |A|$  we have:

$$\sum_{\substack{I+J=A \\ |I|=k}} \prod_{\substack{x \in I \\ y \in J}} \frac{(x - ty/q)(x - y/t)}{(x - y)(x - y/q)} = \sum_{\substack{I+J=A \\ |I|=k}} \prod_{\substack{x \in I \\ y \in J}} \frac{(x - qy/t)(x - ty)}{(x - y)(x - qy)}$$

Firstly make the substitution  $\alpha = t/q$  and  $\beta = 1/t$  to rewrite this as:

$$\sum_{\substack{I+J=A \\ |I|=k}} \prod_{\substack{x \in I \\ y \in J}} \frac{(x - \alpha y)(x - \beta y)}{(x - y)(x - \alpha\beta y)} = \sum_{\substack{I+J=A \\ |I|=k}} \prod_{\substack{x \in I \\ y \in J}} \frac{(\alpha x - y)(\alpha x - y)}{(x - y)(\alpha\beta x - y)}$$

Write  $A = x_1 + x_2 + x_3 + \dots + x_n$ . The  $k = 1$  case reduces to:

$$\sum_{i=1}^n \prod_{j \neq i} \frac{(x_i - \alpha x_j)(x_i - \beta x_j)}{(x_i - x_j)(x_i - \alpha\beta x_j)} = \sum_{i=1}^n \prod_{j \neq i} \frac{(\alpha x_i - x_j)(\beta x_i - x_j)}{(x_i - x_j)(\alpha\beta x_i - x_j)}$$

Now consider the following contour integral:

$$\frac{1}{r!(2\pi i)^k} \oint_{C_1} \dots \oint_{C_r} \frac{dw_1 \dots dw_r}{w_1 \dots w_r} \prod_{i=1}^r \left( \prod_{k=1}^n \frac{(w_i - \alpha x_k)(w_i - \beta x_k)}{(w_i - x_k)(w_i - \alpha\beta x_k)} - 1 \right) \times \prod_{i \neq j} \frac{(w_i - \alpha\beta w_j)(w_i - w_j)}{(w_i - \alpha w_j)(w_i - \beta w_j)}$$

where the contours of integration corresponding to the variables  $w_1, w_2, \dots, w_r$  are taken such that the only poles of the integrand inside these contours are the points  $x_1, \dots, x_n$ . Note that the integrand has no poles at zero or at infinity.

When  $r = 1$  the above integral reduces to:

$$\frac{1}{2\pi i} \oint_C \frac{dw}{w} \left( \prod_{k=1}^n \frac{(w - \alpha x_k)(w - \beta x_k)}{(w - x_k)(w - \alpha\beta x_k)} - 1 \right)$$

By Cauchy's theorem this is equal to the sum over the residues at the poles inside  $C_1$  which occur precisely when  $w = x_i$  for some  $i$ :

$$\frac{(1 - \alpha)(1 - \beta)}{(1 - \alpha\beta)} \left( \sum_i \prod_{j \neq i} \frac{(x_i - \alpha x_j)(x_i - \beta x_j)}{(x_i - x_j)(x_i - \alpha\beta x_j)} \right)$$

One can on the other hand consider the poles outside the contour: they are of the form  $w = \alpha\beta x_i$  for some  $i$ . We obtain:

$$\frac{(\alpha\beta - \alpha)(\alpha\beta - \beta)}{(\alpha\beta - 1)} \left( \sum_i \prod_{j \neq i} \frac{(\alpha\beta x_i - \alpha x_j)(\alpha\beta x_i - \beta x_j)}{\alpha\beta(\alpha\beta x_i - x_j)(\alpha\beta x_i - \alpha\beta x_j)} \right)$$

which simplifies to:

$$-\frac{(1-\alpha)(1-\beta)}{(1-\alpha\beta)} \left( \sum_i \prod_{i \neq j} \frac{(\alpha x_i - x_j)(\beta x_i - x_j)}{(x_i - x_j)(\alpha\beta x_i - x_j)} \right)$$

Noting that the result should be the opposite, we recover the  $k = 1$  version of our identity.

Consider now the case of general  $r$ . We compare once again the residues inside and outside the contours. The integration over each variable  $w_i$  results in a sum over poles of the form  $w_i = x_{a_i}$  for some  $a_i$ . It is easy to check that due to the factors  $w_i - w_j$ , the  $a_i$  must be distinct. Furthermore, due to the symmetry of the integral by exchange of the  $w_i$ , each permutation of the  $a_i$  produces the same contribution and compensates the factor  $r!$ . The result is a sum over subsets  $I = a_1 + \dots + a_r$  and we obtain

$$\begin{aligned} \sum_{\substack{I \subset \{1, \dots, n\} \\ |I|=r}} \frac{1}{x_{a_1} \cdots x_{a_r}} \prod_{i=1}^r \left( \frac{(x_{a_i} - \alpha x_{a_i})(x_{a_i} - \beta x_{a_i})}{(x_{a_i} - \alpha\beta x_{a_i})} \right. \\ \left. \times \prod_{\substack{k=1 \\ k \neq a_i}}^n \frac{(x_{a_i} - \alpha x_k)(x_{a_i} - \beta x_k)}{(x_{a_i} - x_k)(x_{a_i} - \alpha\beta x_k)} \right) \prod_{i \neq j} \frac{(x_{a_i} - x_{a_j})(x_{a_i} - \alpha\beta x_{a_j})}{(x_{a_i} - \alpha x_{a_j})(x_{a_i} - \beta x_{a_j})} \end{aligned}$$

which coincides with the left hand side of our identity after obvious cancellations, up to the factor

$$((1-\alpha)(1-\beta)/(1-\alpha\beta))^r$$

A careful analysis of the poles outside the contours shows that only the poles of the form  $w_i = \alpha\beta x_{a_i}$  contribute, so that we find a similar sum:

$$\begin{aligned} \sum_{\substack{I \subset \{1, \dots, n\} \\ |I|=r}} \frac{1}{(\alpha\beta)^r x_{a_1} \cdots x_{a_r}} \prod_{i=1}^r \left( \frac{(\alpha\beta x_{a_i} - \alpha x_{a_i})(\alpha\beta x_{a_i} - \beta x_{a_i})}{(\alpha\beta x_{a_i} - x_{a_i})} \right) \\ \times \prod_{\substack{k=1 \\ k \neq a_i}}^n \frac{(\alpha\beta x_{a_i} - \alpha x_k)(\alpha\beta x_{a_i} - \beta x_k)}{(\alpha\beta x_{a_i} - x_k)(\alpha\beta x_{a_i} - \alpha\beta x_k)} \prod_{i \neq j} \frac{(\alpha\beta x_{a_i} - x_{a_j})(\alpha\beta x_{a_i} - \alpha\beta x_{a_j})}{(\alpha\beta x_{a_i} - \alpha x_{a_j})(\alpha\beta x_{a_i} - \beta x_{a_j})} \end{aligned}$$

which coincides with the right hand side up to the factor

$$\frac{1}{(\alpha\beta)^r} \left( \frac{(\alpha\beta - \alpha)(\alpha\beta - \beta)}{1 - \alpha\beta} \right)^r$$

□

**Lemma 2.2.**

$$\Phi(z : X) - 1 = \frac{1-t}{1-q} \sum_{x \in X} (w(z : x)\Phi(X - x : x) - W(z : x)\Phi(x : X - x))$$

*Proof.* By the  $k = 1$  case of the previous lemma, both sides vanish as  $z$  goes to infinity, thus one need only compare the residues at the poles.  $\square$

**Proposition 2.1.**

$$\sum_{s=0}^k \frac{(q; t)_s}{(t; t)_s} \sum_{\substack{X' + X'' = X \\ |X'| = k-s}} \left( w(z : X'')V(z : t^{s-1}X'')W(z : t^s X')\Phi(X', X'') \right. \\ \left. - w(z : X')\Phi(X'', X') \right) = 0$$

*Proof.* When  $k = 1$  we have only two terms. When  $s = 1$  we must have  $X' = \emptyset$  and  $X'' = X$  thus the  $s = 1$  term reduces to:

$$\frac{1-q}{1-t} (w(z : X)V(z : X) - 1) = \frac{1-q}{1-t} (\Phi(z : X) - 1)$$

while when  $s = 0$  we must have that  $|X'| = 1$ . Since we have:

$$w(z : X'')V(z : X''/t) = w(z : X'')w(z : X'')^{-1} = 1$$

The  $s = 0$  term becomes:

$$\sum_{x \in X} (W(z : x)\Phi(x : X - x) - w(z : x)\Phi(X - x : x))$$

Thus the  $k = 1$  case of the identity is equivalent to the previous lemma.

More generally, the left hand side is a rational function in  $z$  of degree zero, which vanishes in the limit as  $z$  goes to infinity. The poles are located at:


$$z \in \{a_k t^s, \quad a_k/q : k = 1 \dots m, \quad s = 0 \dots k\}$$

By considering separately the terms for which  $a_k \in X'$  and those for which  $a_k \in X''$  one may check that the residue at the pole  $z = a_k t^s$  vanishes for all  $s$ , while the residue at the pole  $a_k/q$  is equivalent to the  $k - 1$  form of the identity in  $m - 1$  variables.  $\square$


**2.3. Pieri formula and recurrence.**



2.3.1. *Arms and legs again.* For  $\mu \subseteq \lambda$ , let  $C_{\lambda/\mu}$  denote the set of boxes of  $\lambda$  in columns which are longer than the corresponding columns of  $\mu$  and let  $R_{\lambda/\mu}$  denote the set of boxes of  $\lambda$  which are in rows longer than the corresponding rows of  $\mu$ . For example, if  $\lambda = (8, 6, 5, 3, 2)$  and  $\mu = (8, 5, 5, 2, 2)$ ,



Let  $\tilde{C}_{\lambda/\mu}$  denote the set of boxes of  $\lambda$  in columns which are the same length as the corresponding columns of  $\mu$ , and let  $\tilde{R}_{\lambda/\mu}$  denote the set of boxes of  $\lambda$  in rows which are the same length as the corresponding rows of  $\mu$ . With the same example,



Now let us define:

$$\begin{aligned}
 C_{\lambda/\mu}(q, t) &= \sum_{s \in C_{\lambda/\mu}} q^{a_\lambda(s)} t^{l_\lambda(s)} - \sum_{s \in \mu \cap C_{\lambda/\mu}} q^{a_\mu(s)} t^{l_\mu(s)} \\
 R_{\lambda/\mu}(q, t) &= \sum_{s \in R_{\lambda/\mu}} q^{a_\lambda(s)} t^{l_\lambda(s)} - \sum_{s \in \mu \cap R_{\lambda/\mu}} q^{a_\mu(s)} t^{l_\mu(s)} \\
 \tilde{C}_{\lambda/\mu}(q, t) &= \sum_{s \in \tilde{C}_{\lambda/\mu}} (q^{a_\lambda(s)} t^{l_\lambda(s)} - q^{a_\mu(s)} t^{l_\mu(s)}) \\
 \tilde{R}_{\lambda/\mu}(q, t) &= \sum_{s \in \tilde{R}_{\lambda/\mu}} (q^{a_\lambda(s)} t^{l_\lambda(s)} - q^{a_\mu(s)} t^{l_\mu(s)})
 \end{aligned}$$

It is clear that we have:

$$\begin{aligned}
 C_{\lambda'/\mu'}(t, q) &= R_{\lambda/\mu}(q, t) \\
 \tilde{C}_{\lambda'/\mu'}(t, q) &= \tilde{R}_{\lambda/\mu}(q, t)
 \end{aligned}$$

as well as that:

$$\begin{aligned}
 B_\lambda(q, t) - B_\mu(q, t) &= C_{\lambda/\mu}(q, t) + \tilde{C}_{\lambda/\mu}(q, t) \\
 &= R_{\lambda/\mu}(q, t) + \tilde{R}_{\lambda/\mu}(q, t)
 \end{aligned}$$

2.3.2. *Pieri formula.* The Macdonald polynomials are, in fact, the eigenfunctions of the more general family of operators [29] (page 315):

$$\frac{1}{\Delta_n(X)} \sum_{\substack{I \subseteq [n] \\ |I|=r}} \Delta_n(X + (1-t)X_I) f(X + (1-q)X_I)$$

where  $X_I$  is the alphabet:

$$X_I = \sum_{i \in I} x_i$$

with eigenvalues:

$$e_r \left( \sum_{i=1}^n q^{\lambda_i} t^{n-i} \right)$$

Using this fact, Macdonald was also able to show that his polynomials satisfy the following generalization of the Pieri formulae [29] (page 340):

$$\begin{aligned} P_\mu(X) \Omega_z \left( X \frac{1-t}{1-q} \right) &= \sum_{\lambda \in U(\mu)} \varphi_{\lambda/\mu}(q, t) P_\lambda(q, t) z^{|\lambda| - |\mu|} \\ Q_\mu(X) \tilde{\Omega}_z(X) &= \sum_{\lambda \in \tilde{U}(\mu)} \varphi'_{\lambda/\mu}(q, t) Q_\lambda(X) z^{|\lambda| - |\mu|} \end{aligned}$$

where:

$$\begin{aligned} \varphi_{\lambda/\mu}(q, t) &= \Omega((q-t)C_{\lambda/\mu}(q, t)) \\ \varphi'_{\lambda/\mu}(q, t) &= \Omega((t-q)R_{\lambda/\mu}(q, t)) \end{aligned}$$

As before  $U(\mu)$  denotes the set of partitions obtained from  $\mu$  by adding a horizontal strip while  $\tilde{U}(\mu)$  denotes the set of partitions obtained from  $\mu$  by adding a vertical strip. Note that the latter may be obtained from the former by applying the operator  $\omega_{q,t}$  to both sides and then formally replacing  $\lambda$  by  $\lambda'$  and  $\mu$  by  $\mu'$ . The two additional Pieri formulae:

$$\begin{aligned} Q_\mu(X) \Omega_z \left( X \frac{1-t}{1-q} \right) &= \sum_{\lambda \in U(\mu)} \psi_{\lambda/\mu}(q, t) Q_\lambda(X) z^{|\lambda| - |\mu|} \\ P_\mu(X) \tilde{\Omega}_z(X) &= \sum_{\lambda \in \tilde{U}(\mu)} \psi'_{\lambda/\mu}(q, t) P_\lambda(X) z^{|\lambda| - |\mu|} \end{aligned}$$

where:

$$\begin{aligned}\psi_{\lambda/\mu}(q, t) &= \Omega((t - q)\tilde{C}_{\lambda/\mu}(q, t)) \\ \psi'_{\lambda/\mu}(q, t) &= \Omega((q - t)\tilde{R}_{\lambda/\mu}(q, t))\end{aligned}$$

are obtained from by multiplying, or dividing by  $\Omega((q - t)(B_\lambda(q, t) - B_\mu(q, t)))$  as appropriate.

2.3.3. *Recurrence.* Dual to the Pieri formulae, we have the following recurrences [29] (pages 346, 348):

$$\begin{aligned}P_\lambda(X + z) &= \sum_{\mu \in D(\lambda)} \psi_{\lambda/\mu}(q, t) P_\mu(X) z^{|\lambda| - |\mu|} \\ Q_\lambda(X - z) &= \sum_{\mu \in \tilde{D}(\lambda)} \psi'_{\lambda/\mu}(q, t) Q_\mu(X) (\epsilon z)^{|\lambda| - |\mu|} \\ Q_\lambda(X + z) &= \sum_{\mu \in D(\lambda)} \varphi_{\lambda/\mu}(q, t) Q_\mu(X) z^{|\lambda| - |\mu|} \\ P_\lambda(X - z) &= \sum_{\mu \in \tilde{D}(\lambda)} \varphi'_{\lambda/\mu}(q, t) P_\mu(X) (\epsilon z)^{|\lambda| - |\mu|}\end{aligned}$$

Here  $D(\lambda)$  denotes the set of partitions obtained from  $\lambda$  by removing a horizontal strip, while  $\tilde{D}(\lambda)$  denotes the set of partitions obtained from  $\lambda$  by removing a vertical strip.

Note that it is also possible to take any one of the recurrences or the Pieri formulae, together with the definition of the Macdonald polynomial of a row, or a column accordingly as the *definition* of the Macdonald polynomials.

## 2.4. The Proof.

2.4.1. *The Schur case.* To aid the reader in not getting lost in a thicket of  $q$ 's and  $t$ 's we first give a proof of the classical Schur identity, which mirrors, in simplified form, the main steps of the proof of the more complicated Kawanaka identity.

### Proposition 2.2.

$$\sum_{\lambda} s_{\lambda}(x_1, \dots, x_{n+1}) = \prod_{i=1}^n \frac{1}{(1 - x_i)} \prod_{1 \leq i < j \leq n} \frac{1}{(1 - x_i x_j)}$$

*Proof.* The proof is by induction on the number of variables. The first step makes use of the Pieri formula and recurrence for Schur functions. The second step is a bijection between two families of partitions.

When  $n = 1$  we get the expansion of the geometric series:

$$\sum_{k \geq 0} x_1^k = \frac{1}{1 - x_1}$$

By the recurrence for the Schur functions we have:

$$\begin{aligned} \text{LHS}(n+1) &= \sum_{\lambda} s_{\lambda}(x_1, \dots, x_{n+1}) \\ &= \sum_{\lambda} \sum_{\mu \in D(\lambda)} s_{\mu}(x_1, \dots, x_n) x_{n+1}^{|\lambda| - |\mu|} \\ &= \sum_{\mu} s_{\mu}(x_1, \dots, x_n) \sum_{\lambda \in U(\mu)} x_{n+1}^{|\lambda| - |\mu|} \end{aligned}$$

By the induction hypothesis:

$$\begin{aligned} \text{RHS}(n+1) &= \prod_{i=1}^{n+1} \frac{1}{1 - x_i} \prod_{1 \leq i < j \leq n+1} \frac{1}{1 - x_i x_j} \\ &= \frac{1}{1 - x_{n+1}} \prod_{i=1}^n \frac{1}{1 - x_i} \prod_{1 \leq i < j \leq n} \frac{1}{1 - x_i x_j} \prod_{i=1}^n \frac{1}{1 - x_i x_{n+1}} \\ &= \frac{1}{1 - x_{n+1}} \text{RHS}(n) \prod_{i=1}^n \frac{1}{1 - x_i x_{n+1}} \\ &= \frac{1}{1 - x_{n+1}} \text{LHS}(n) \prod_{i=1}^n \frac{1}{1 - x_i x_{n+1}} \\ &= \frac{1}{1 - x_{n+1}} \sum_{\mu} s_{\mu}(x_1, \dots, x_n) \prod_{i=1}^n \frac{1}{1 - x_i x_{n+1}} \end{aligned}$$

But by the Pieri formula for the Schur functions:

$$\begin{aligned} \sum_{\mu} s_{\mu}(x_1, \dots, x_n) \prod_{i=1}^n \frac{1}{1 - x_i x_{n+1}} &= \sum_{\mu} s_{\mu}(x_1, \dots, x_n) \sum_s h_s(x_1, \dots, x_n) x_{n+1}^s \\ &= \sum_{\mu} \sum_{\lambda \in U(\mu)} s_{\lambda}(x_1, \dots, x_n) x_{n+1}^{|\lambda| - |\mu|} \\ &= \sum_{\lambda} s_{\lambda}(x_1, \dots, x_n) \sum_{\mu \in D(\lambda)} x_{n+1}^{|\lambda| - |\mu|} \end{aligned}$$

By considering the coefficient of  $s_\mu(X)$  on both sides, it suffices to demonstrate that:

$$(12) \quad \sum_{\lambda \in U(\mu)} x_{n+1}^{|\lambda| - |\mu|} = \frac{1}{1 - x_{n+1}} \sum_{\gamma \in D(\mu)} x_{n+1}^{|\mu| - |\gamma|}$$

Conjugating all the partitions involved, this is equivalent to:

$$(13) \quad \sum_{\lambda \in \tilde{U}(\mu)} x_{n+1}^{|\lambda| - |\mu|} = \frac{1}{1 - x_{n+1}} \sum_{\gamma \in \tilde{D}(\mu)} x_{n+1}^{|\mu| - |\gamma|}$$


Suppose now that  $\mu$  is a partition with distinct  $m$  distinct nonzero parts. For each  $\alpha \subseteq [m]$  such that  $|\alpha| = k$  let  $\mu_-(\alpha)$  denote the partition obtained from  $\mu$  by removing a box from the end of each row indexed by an element of  $\alpha$ . We have:

$$\tilde{D}_k(\mu) = \{\mu_-(\alpha), \alpha \subseteq [m], |\alpha| = k\}$$

Similarly, let  $\mu_+(\alpha)$  denote the partition obtained from  $\mu$  by adding a box to the end of each row indexed by  $\alpha$ . Let:

$$S\tilde{U}_k(\mu) = \{\mu_+(\alpha), \alpha \subseteq [m], |\alpha| = k\}$$

There is a natural bijection between  $\tilde{D}_k(\mu)$  and  $S\tilde{U}_k(\mu)$  which send  $\mu_-(\alpha)$  to  $\mu_+(\alpha)$ .



If  $\mu$  contains repeated parts, then it is no longer true that  $\mu_+(\alpha)$  and  $\mu_-(\alpha)$  will be valid partitions for all  $\alpha$ . For example, if  $\mu_i = \mu_{i+1}$  and  $i \in \alpha$  but  $(i+1) \notin \alpha$  then  $\mu_-(\alpha)$  will not be a valid partition.

Conversely, if  $i \notin \alpha$  while  $(i+1) \in \alpha$  then  $\mu_+(\alpha)$  will not be a valid partition. There is still, however a natural bijection between  $\tilde{D}_k(\mu)$  and  $S\tilde{U}_k(\mu)$ , though it is slightly more cumbersome to describe.

More generally for each  $\alpha \subseteq [m]$  such that  $|\alpha| = k$  and for each  $p \geq 0$  let  $\mu_+(\alpha, p)$  denote the partition obtained from  $\mu$  by adding a box to the end of each row indexed by an element of  $\alpha$ , and then adding  $p$  new rows each of length one. Let:

$$S\tilde{U}_k(\mu, p) = \{\mu_+(\alpha, p), \alpha \subseteq [m], |\alpha| = k\}$$

Upon consideration of the coefficient of  $x_n^k$  in equation (13), to complete the proof we must show that:

$$(14) \quad |\tilde{U}_k(\mu)| = \sum_s |\tilde{D}_{k-s}(\mu)|$$

This follows immediately from the fact that:

$$\tilde{U}_k(\mu) = \bigcup_{p=0}^k S\tilde{U}_{k-p}(\mu, p)$$

and that for each  $p$  we have:

$$|S\tilde{U}_k(\mu, p)| = |S\tilde{U}_k(\mu, 0)| = |\tilde{D}_k(\mu)|$$

□

2.4.2. *Step one.* For the Kawanaka identity, let us define:

$$\begin{aligned} \text{LHS}(n) &= \sum_{\lambda} \Omega((q - \epsilon t) B_{\lambda}(q, t)) P_{\lambda}(X_n; q^2, t^2) \\ \text{RHS}(n) &= \Omega\left(\frac{1 - \epsilon t}{1 - q} X_n + \frac{1 - t^2}{1 - q^2} e_2(X_n)\right) \end{aligned}$$

The case  $n = 1$  is a special case of the  $q$ -binomial formula:

$$\begin{aligned} \Omega\left(\frac{1 - \epsilon t}{1 - q} x_1\right) &= \sum_n \Omega\left(\frac{(q - \epsilon t)(1 - q^n)}{(1 - q)}\right) x_1^n \\ &= \sum_n \Omega((q - \epsilon t) B_{(n)}(q, \epsilon t)) x_1^n \end{aligned}$$

We have now, by the recurrence for the Macdonald  $P$ -functions:

$$\begin{aligned} &\text{LHS}(n+1) \\ &= \sum_{\lambda} \Omega((q - \epsilon t) B_{\lambda}(q, t)) P_{\lambda}(X_n + x_{n+1}; q^2, t^2) \\ &= \sum_{\lambda} \Omega((q - \epsilon t) B_{\lambda}(q, t)) \sum_{\mu \in D(\lambda)} \Omega((t^2 - q^2) \tilde{C}_{\lambda/\mu}(q^2, t^2)) P_{\mu}(X_n; q^2, t^2) x_{n+1}^{|\lambda| - |\mu|} \\ &= \sum_{\mu} P_{\mu}(X_n, q^2, t^2) \sum_k x_{n+1}^k \sum_{\lambda \in U_k(\mu)} \Omega((q - \epsilon t) B_{\lambda}(q, t)) \Omega((t^2 - q^2) \tilde{C}_{\lambda/\mu}(q^2, t^2)) \end{aligned}$$

while, by the induction hypothesis:

$$\begin{aligned} \text{RHS}(n+1) &= \Omega\left(\frac{1-\epsilon t}{1-q}x_{n+1}\right) \text{RHS}(n) \Omega\left(\frac{1-t^2}{1-q^2}X_n x_{n+1}\right) \\ &= \Omega\left(\frac{1-\epsilon t}{1-q}x_{n+1}\right) \text{LHS}(n) \Omega\left(\frac{1-t^2}{1-q^2}X_n x_{n+1}\right) \end{aligned}$$

But, by the Pieri formula for the Macdonald  $P$  functions, we have:

$$\begin{aligned} &\text{LHS}(n) \Omega\left(\frac{1-t^2}{1-q^2}X_n x_{n+1}\right) \\ &= \sum_{\gamma} \Omega((q-\epsilon t)B_{\gamma}(q, t)) P_{\gamma}(X_n; q^2, t^2) \Omega\left(\frac{1-t^2}{1-q^2}X_n x_{n+1}\right) \\ &= \sum_{\gamma} \Omega((q-\epsilon t)B_{\gamma}(q, t)) \sum_{\mu \in U(\gamma)} \Omega((q^2-t^2)C_{\mu/\gamma}(q^2, t^2)) P_{\mu}(X_n; q^2, t^2) x_{n+1}^{|\mu|-|\gamma|} \\ &= \sum_{\mu} P_{\mu}(X_n; q^2, t^2) \sum_k x_{n+1}^k \sum_{\gamma \in D_k(\mu)} \Omega((q-\epsilon t)B_{\gamma}(q, t)) \Omega((q^2-t^2)C_{\mu/\gamma}(q^2, t^2)) \end{aligned}$$

By considering the coefficient of  $P_{\mu}(X; q^2, t^2)$  on both sides, and then dividing by  $\Omega((q-\epsilon t)B_{\mu}(q, t))$ , we need to show that, for any  $\mu$  we have:

$$\begin{aligned} &\sum_k x_{n+1}^k \sum_{\lambda \in U_k(\mu)} \Omega((q-\epsilon t)(B_{\lambda}(q, t) - B_{\mu}(q, t))) \Omega((t^2-q^2)\tilde{C}_{\lambda/\mu}(q^2, t^2)) \\ &= \Omega\left(\frac{1-\epsilon t}{1-q}x_{n+1}\right) \times \\ &\sum_k x_{n+1}^k \sum_{\gamma \in D_k(\mu)} \Omega((q-\epsilon t)(B_{\gamma}(q, t) - B_{\mu}(q, t))) \Omega((q^2-t^2)C_{\mu/\gamma}(q^2, t^2)) \end{aligned}$$

Observe that when  $q = \epsilon t$  this reduces to equation (12).

By conjugating all the partitions involved, and interchanging the roles of  $q$  and  $t$  we obtain the dual form:

$$\begin{aligned} &\sum_k x_{n+1}^k \sum_{\lambda \in \tilde{U}_r(\mu)} \Omega((t-\epsilon q)(B_{\lambda}(q, t) - B_{\mu}(q, t))) \Omega((q^2-t^2)\tilde{R}_{\lambda/\mu}(q^2, t^2)) \\ &= \Omega\left(\frac{1-\epsilon q}{1-t}x_{n+1}\right) \times \\ &\sum_k x_{n+1}^k \sum_{\gamma \in \tilde{D}_z(\mu)} \Omega((t-\epsilon q)(B_{\gamma}(q, t) - B_{\mu}(q, t))) \Omega((t^2-q^2)R_{\lambda/\mu}(q^2, t^2)) \end{aligned}$$

Here we have made use of the fact that  $B_{\lambda'}(t, q) = B_{\lambda}(q, t)$  as well as  $\tilde{C}_{\lambda'/\mu'}(t, q) = \tilde{R}_{\lambda/\mu}(q, t)$  and  $C_{\lambda'/\mu'}(t, q) = R_{\lambda/\mu}(q, t)$ .

Note that when  $q = \epsilon t$  this reduces to equation (13).

Finally, making use of the  $q$ -binomial formula once again, on consideration of the coefficient of  $x_{n+1}^r$  on both sides, we must show that for any  $\mu$  and any  $r$  we have:

$$\begin{aligned}
 (15) \quad & \sum_{\lambda \in \tilde{U}_r(\mu)} \Omega((t - \epsilon q)(B_{\lambda}(q, t) - B_{\mu}(q, t))) \Omega((q^2 - t^2)R_{\lambda/\mu}(q^2, t^2)) \\
 &= \sum_s \Omega((t - \epsilon q)B_{(1^r-s)}(\epsilon q, t)) \times \\
 & \sum_{\gamma \in \tilde{D}_s(\mu)} \Omega((t - \epsilon q)(B_{\gamma}(q, t) - B_{\mu}(q, t))) \Omega((t^2 - q^2)\tilde{R}_{\lambda/\mu}(q^2, t^2))
 \end{aligned}$$

Let us define:

$$\begin{aligned}
 H_{\lambda}(s) &= \Omega((q - \epsilon t)q^{a_{\lambda}(s)}t^{l_{\lambda}(s)}) = \frac{1 + q^{a_{\lambda}(s)}t^{l_{\lambda}(s)+1}}{1 - q^{a_{\lambda}(s)+1}t^{l_{\lambda}(s)}} \\
 \tilde{H}_{\lambda}(s) &= \Omega((t - \epsilon q)q^{a_{\lambda}(s)}t^{l_{\lambda}(s)}) = \frac{1 + q^{a_{\lambda}(s)+1}t^{l_{\lambda}(s)}}{1 - q^{a_{\lambda}(s)}t^{l_{\lambda}(s)+1}} \\
 G_{\lambda}(s) &= \Omega((t - q)q^{2a_{\lambda}(s)}t^{2l_{\lambda}(s)}) = \frac{1 - q^{2(a_{\lambda}(s)+1)}t^{2l_{\lambda}(s)}}{1 - q^{2a_{\lambda}(s)}t^{2(l_{\lambda}(s)+1)}}
 \end{aligned}$$

We have, by the difference of perfect squares  $\Omega(a^2) = \Omega(a + \epsilon a)$  that:

$$\begin{aligned}
 G_{\lambda}(s)H_{\lambda}(s) &= \Omega((t^2 - q^2)q^{2a_{\lambda}(s)}t^{2l_{\lambda}(s)} + (q - \epsilon t)q^{a_{\lambda}(s)}t^{l_{\lambda}(s)}) \\
 &= \Omega((t - \epsilon q)q^{a_{\lambda}(s)}t^{l_{\lambda}(s)}) \\
 &= \tilde{H}_{\lambda}(s)
 \end{aligned}$$



Now we may simplify equation (15) by writing:

$$\begin{aligned}
 L(\lambda, \mu) &= \Omega((t - \epsilon q)(B_\lambda(q, t) - B_\mu(q, t))) \Omega((q^2 - t^2) \tilde{R}_{\lambda/\mu}(q^2, t^2)) \\
 &= \Omega\left((q - \epsilon t) \tilde{R}_{\lambda/\mu}(q, t) + (t - \epsilon q) R_{\lambda/\mu}(q, t)\right) \\
 &= \prod_{s \in \tilde{R}_{\lambda/\mu}} \frac{H_\lambda(s)}{H_\mu(s)} \prod_{s \in R_{\lambda/\mu}} \frac{\tilde{H}_\lambda(s)}{\tilde{H}_\mu(s)} \\
 &= \prod_{s \in \tilde{R}_{\lambda/\mu} \cap C_{\lambda/\mu}} \frac{H_\lambda(s)}{H_\mu(s)} \prod_{s \in R_{\lambda/\mu}} \frac{\tilde{H}_\lambda(s)}{\tilde{H}_\mu(s)}
 \end{aligned}$$

and:

$$\begin{aligned}
 R(\mu, \gamma) &= \Omega((t - \epsilon q)(B_\gamma(q, t) - B_\mu(q, t))) \Omega((t^2 - q^2) \tilde{R}_{\mu/\gamma}(q^2, t^2)) \\
 &= \Omega\left(-(q - \epsilon t) R_{\mu/\gamma}(q, t) - (t - \epsilon q) \tilde{R}_{\mu/\gamma}(q, t)\right) \\
 &= \prod_{s \in R_{\mu/\gamma}} \frac{H_\gamma(s)}{H_\mu(s)} \prod_{s \in \tilde{R}_{\mu/\gamma}} \frac{\tilde{H}_\gamma(s)}{\tilde{H}_\mu(s)} \\
 &= \prod_{s \in R_{\mu/\gamma}} \frac{H_\gamma(s)}{H_\mu(s)} \prod_{s \in \tilde{R}_{\mu/\gamma} \cap C_{\mu/\gamma}} \frac{\tilde{H}_\gamma(s)}{\tilde{H}_\mu(s)}
 \end{aligned}$$

To complete the proof we must show that for any partition  $\mu$  and for any positive integer  $k$  we have:

$$(16) \quad \sum_{\lambda \in \tilde{U}_k(\mu)} L(\lambda, \mu) = \sum_s \frac{(-q; t)_s}{(t; t)_s} \sum_{\gamma \in \tilde{D}_{k-s}(\mu)} R(\mu, \gamma)$$

Note that when  $q = \epsilon t$  we have  $L(\lambda, \mu) = R(\mu, \gamma) = 1$  and the above reduces to equation (14).

**2.4.3. Step two.** Suppose that  $\mu$  is a partition with  $m$  distinct nonzero parts. For each  $k \in [m] = \{1, 2, \dots, m\}$  let  $a_k = q^{\mu_k} t^{m-k}$ . For any  $K \subseteq [m]$  let  $A_K$  denote the alphabet:

$$A_K = \sum_{k \in K} a_k$$

The next step is to obtain several related combinatorial descriptions for the rational function first introduced in section 2.2:


$$\Phi(A_I, A_J)_{\epsilon q, t} = \prod_{\substack{i \in I \\ j \in J}} \frac{(a_i + t/q \ a_j)(a_i - a_j/t)}{(a_i - a_j)(a_i + a_j/q)}$$

Note that  $q$  has been replaced by  $\epsilon q$ , that is every occurrence of  $q$  has been replaced by its formal (as opposed to plethystic) negative.

Let  $\alpha$  and  $\tilde{\alpha}$  be such that  $\alpha \cup \tilde{\alpha} = [m]$ . Let  $\gamma = \mu_-(\alpha)$  denote the partition obtained from  $\mu$  by removing a box from the end of each row indexed by  $\alpha$  and let  $\tilde{\gamma} = \mu_-(\tilde{\alpha})$  denote the partition obtained from  $\mu$  by removing a box from the end of each row indexed by  $\tilde{\alpha}$ .

If  $s \in \tilde{R}_{\mu/\gamma} \cap C_{\mu/\gamma}$  then  $s$  must be of the form  $s = (i, \mu_j)$  for some  $j \in \alpha$  and  $i \notin \alpha$  with  $i < j$ . In this case we have that  $a_\gamma(s) = a_\mu(s) = \mu_i - \mu_j$ , while  $l_\gamma(s) + 1 = l_\mu(s) = j - i$ .

For example:



$\mu = (8, 6, 5, 3, 2)$  and  $\alpha = \{2, 4\}$ ,  $\tilde{R}_{\mu/\gamma} \cap C_{\mu/\gamma} = \{(1, \mu_4), (1, \mu_2), (3, \mu_4)\} = \{(1, 3), (1, 6), (3, 3)\}$ .

In this case, one may check that:

$$\tilde{H}_\mu(i, \mu_j) = \frac{1 + q a_i a_j^{-1}}{1 - t a_i a_j^{-1}} = v(a_j : a_i)_{\epsilon q, t}^{-1}$$


$$\tilde{H}_{\mu_-(\alpha)}(i, \mu_j) = \frac{1 + q/t a_i a_j^{-1}}{1 - a_i a_j^{-1}} = W(a_j : a_i)_{\epsilon q, t}$$

so that:

$$(17) \quad \frac{\tilde{H}_{\mu_-(\alpha)}(i, \mu_j)}{\tilde{H}_\mu(i, \mu_j)} = \Theta(a_j, a_i)_{\epsilon q, t} = \Phi(a_i, a_j)_{\epsilon q, t}$$

Similarly if  $s \in \tilde{R}_{\mu/\tilde{\gamma}} \cap C_{\mu/\tilde{\gamma}}$  then  $s$  must be of the form  $s = (j, \mu_i)$  for some  $j \notin \tilde{\alpha}$  and  $i \in \tilde{\alpha}$  with  $j < i$ . now  $a_{\tilde{\gamma}}(s) = a_\mu(s) = \mu_j - \mu_i$ , while  $l_{\tilde{\gamma}}(s) + 1 = l_\mu(s) = i - j$ .

For example:



$\mu = (8, 6, 5, 3, 2)$  and  $\tilde{\alpha} = \{1, 3, 5\}$ ,  $\tilde{R}_{\mu/\tilde{\gamma}} \cap C_{\mu/\tilde{\gamma}} = \{(2, \mu_5), (2, \mu_3), (4, \mu_5)\} = \{(2, 2), (2, 5), (4, 2)\}$ .

In this case we have:

$$\begin{aligned}\tilde{H}_{\mu}(j, \mu_i) &= v(a_i : a_j)_{\epsilon q, t}^{-1} \\ \tilde{H}_{\mu_-(\tilde{\alpha})}(j, \mu_i) &= W(a_i : a_j)_{\epsilon q, t}\end{aligned}$$


so that:

$$(18) \quad \frac{\tilde{H}_{\mu_-(\tilde{\alpha})}(j, \mu_i)}{\tilde{H}_{\mu}(j, \mu_i)} = \Theta(a_i, a_j)_{\epsilon q, t} = \Phi(a_j, a_i)_{\epsilon q, t}$$

Now let  $\lambda = \mu_+(\alpha)$  and let  $\tilde{\lambda} = \mu_+(\tilde{\alpha})$ .

If  $s \in \tilde{R}_{\lambda/\mu} \cap C_{\lambda/\mu}$  then we must have  $s = (i, \mu_j + 1)$  for some  $j \in \alpha$  and  $i \notin \alpha$  with  $i < j$ . This time we have that  $a_{\mu}(s) = a_{\lambda}(s) = \mu_i - \mu_j - 1$ , while  $l_{\mu}(s) + 1 = l_{\lambda}(s) = j - i$ .

For example:



If  $\mu = (8, 6, 5, 3, 2)$  and  $\alpha = \{2, 4\}$  then  $\tilde{R}_{\lambda/\mu} \cap C_{\lambda/\mu} = \{(1, \mu_4 + 1), (1, \mu_2 + 1), (3, \mu_4 + 1)\} = \{(1, 4), (1, 7), (3, 4)\}$ .

Now:


$$\begin{aligned}H_{\mu_+(\alpha)}(i, \mu_j + 1) &= \frac{1 + t/q \, a_i a_j^{-1}}{1 - a_i a_j^{-1}} = V(a_j, a_i)_{\epsilon q, t} \\ H_{\mu}(i, \mu_j + 1) &= \frac{1 + 1/q \, a_i a_j^{-1}}{1 - 1/t \, a_i a_j^{-1}} = w(a_j : a_i)_{\epsilon q, t}^{-1}\end{aligned}$$

so that:

$$(19) \quad \frac{H_{\mu_+(\alpha)}(i, \mu_j + 1)}{H_{\mu}(i, \mu_j + 1)} = \Phi(a_j, a_i)_{\epsilon q, t}$$

Similarly, if  $s \in \tilde{R}_{\tilde{\lambda}/\mu} \cap C_{\tilde{\lambda}/\mu}$  then we must have  $s = (j, \mu_i + 1)$  for some  $j \notin \tilde{\alpha}$  and  $i \in \tilde{\alpha}$  with  $j < i$ . This time  $a_{\mu}(s) = a_{\lambda}(s) = \mu_j - \mu_i - 1$ , while  $l_{\mu}(s) + 1 = l_{\lambda}(s) = i - j$ .

For example:



If  $\mu = (8, 6, 5, 3, 2)$  and  $\tilde{\alpha} = \{1, 3, 5\}$  then  $\tilde{R}_{\tilde{\lambda}/\mu} \cap C_{\tilde{\lambda}/\mu} = \{(2, \mu_5 + 1), (2, \mu_3 + 1), (4, \mu_5 + 1)\} = \{(2, 3), (2, 6), (4, 3)\}$

We have:

$$\begin{aligned} H_{\mu_+(\tilde{\alpha})}(j, \mu_i + 1) &= V(a_i, a_j)_{\epsilon q, t} \\ H_{\mu}(j, \mu_i + 1) &= w(a_i, a_j)_{\epsilon q, t}^{-1} \end{aligned}$$

so that:

$$(20) \quad \frac{H_{\mu_+(\tilde{\alpha})}(j, \mu_i + 1)}{H_{\mu}(j, \mu_i + 1)} = \Phi(a_i, a_j)_{\epsilon q, t}$$

Note that in equations (17) and (19) we have the restriction that  $i < j$  while in equations (18) and (20) we have the restriction that  $j < i$ .

We may combine all these observations into the following:

**Proposition 2.3.** *We have:*


$$\Phi(A_I, A_J)_{\epsilon q, t} = \prod_{\substack{i \notin \alpha \\ j \in \alpha \\ i < j}} \frac{\tilde{H}_{\mu_-(\alpha)}(i, \mu_j)}{\tilde{H}_{\mu}(i, \mu_j)} \prod_{\substack{i \notin \alpha \\ j \in \alpha \\ j < i}} \frac{H_{\mu_+(\tilde{\alpha})}(j, \mu_i + 1)}{H_{\mu}(j, \mu_i + 1)}$$

and:

$$\Phi(A_J, A_I)_{\epsilon q, t} = \prod_{\substack{i \notin \alpha \\ j \in \alpha \\ i < j}} \frac{H_{\mu_+(\alpha)}(i, \mu_j + 1)}{H_{\mu}(i, \mu_j + 1)} \prod_{\substack{i \notin \alpha \\ j \in \alpha \\ j < i}} \frac{\tilde{H}_{\mu_-(\tilde{\alpha})}(j, \mu_i)}{\tilde{H}_{\mu}(j, \mu_i)}$$

*Proof.* The first expression is obtained by combining equations (17) and (20). The second expression is obtained by combining equations (18) and (19).  $\square$

Now let  $\underline{\mu}$  be the partition obtained from  $\mu$  by removing a box to the end of *every* row (or equivalently removing the first column), and let  $\overline{\mu}$  be the partition obtained from  $\mu$  by *adding* a box to every row.



Although the highlighted box has the same arm and leg length in every partition, it has different co-ordinates with respect to the top left hand corner, giving us:

$$H_{\underline{\mu}}(i-1, \mu_j) = H_{\mu}(i, \mu_j) = H_{\overline{\mu}}(i+1, \mu_j+1)$$

and similarly:

$$\tilde{H}_{\underline{\mu}}(i-1, \mu_j) = \tilde{H}_{\mu}(i, \mu_j) = \tilde{H}_{\overline{\mu}}(i+1, \mu_j+1)$$

Observe further that  $\overline{\mu}_-(\tilde{\alpha}) = \mu_+(\alpha)$  while  $\underline{\mu}_+(\tilde{\alpha}) = \mu_-(\alpha)$ . Using these facts we may rewrite the previous proposition as:

**Proposition 2.4.** *We have:*

$$\Phi(A_I, A_J)_{\epsilon q, t} = \prod_{\substack{i \notin \alpha \\ j \in \alpha \\ i < j}} \frac{\tilde{H}_{\mu_-(\alpha)}(i, \mu_j)}{\tilde{H}_{\mu}(i, \mu_j)} \prod_{\substack{i \notin \alpha \\ j \in \alpha \\ j < i}} \frac{H_{\mu_-(\alpha)}(j, \mu_i)}{H_{\mu}(j, \mu_i + 1)}$$

and:

$$\Phi(A_J, A_I)_{\epsilon q, t} = \prod_{\substack{i \notin \alpha \\ j \in \alpha \\ i < j}} \frac{H_{\mu_+(\alpha)}(i, \mu_j + 1)}{H_{\mu}(i, \mu_j + 1)} \prod_{\substack{i \notin \alpha \\ j \in \alpha \\ j < i}} \frac{\tilde{H}_{\mu_+(\alpha)}(j, \mu_i + 1)}{\tilde{H}_{\mu}(j, \mu_i)}$$

*Proof.* This proposition reduces to the previous one after observing in the first case that:

$$H_{\mu_-(\alpha)}(j, \mu_i) = H_{\underline{\mu}_+(\bar{\alpha})}(j, \underline{\mu}_i + 1) = H_{\mu_+(\bar{\alpha})}(j, \mu_i + 1)$$

and in the second case that:

$$\tilde{H}_{\mu_+(\alpha)}(j, \mu_i + 1) = \tilde{H}_{\bar{\mu}_-(\bar{\alpha})}(j, \bar{\mu}_i) = \tilde{H}_{\mu_-(\bar{\alpha})}(j, \mu_i)$$

□

2.4.4. *Step three.* Using proposition 2.4 we may now give an alternative description for the expressions  $R(\mu, \gamma)$  and  $L(\mu, \gamma)$  arising in equation (16).

**Proposition 2.5.** *If  $\gamma = \mu_-(\alpha)$  then:*

$$R(\mu, \gamma) = w(1/t : A_J)_{\epsilon q, t} \Phi(A_I, A_J)_{\epsilon q, t}$$


*Proof.* By equation (17) we can write:

$$\begin{aligned} \prod_{s \in \tilde{R}_{\mu/\gamma} \cap C_{\mu/\gamma}} \frac{\tilde{H}_{\gamma}(s)}{\tilde{H}_{\mu}(s)} &= \prod_{\substack{i \notin \alpha \\ j \in \alpha \\ i < j}} \frac{\tilde{H}_{\mu_-(\alpha)}(i, \mu_j)}{\tilde{H}_{\mu}(i, \mu_j)} \\ &= \prod_{\substack{i \notin \alpha \\ j \in \alpha \\ i < j}} \Phi(a_i, a_j)_{\epsilon q, t} \end{aligned}$$

On the other hand, by shifting indices, we have:

$$\begin{aligned} \prod_{s \in R_{\mu/\gamma}} \frac{H_{\gamma}(s)}{H_{\mu}(s)} &= \prod_{j \in \alpha} \left( \prod_{k=1}^{\mu_j-1} \frac{H_{\mu_-(\alpha)}(j, k)}{H_{\mu}(j, k)} \right) \frac{1}{H_{\mu}(j, \mu_j)} \\ &= \prod_{j \in \alpha} \frac{1}{H_{\mu}(j, 1)} \left( \prod_{k=1}^{\mu_j-1} \frac{H_{\mu_-(\alpha)}(j, k)}{H_{\mu}(j, k+1)} \right) \end{aligned}$$

Suppose that we take the diagram for  $\gamma = \mu_-(\alpha)$  and place it on top of the diagram for  $\mu$ , but shifted to the right by one column.



For  $j \in \alpha$ , we have that  $H_\mu(j, k+1) = H_{\mu_-(\alpha)}(j, k)$  unless  $k = \mu_i$  for some  $i \notin \alpha$  with  $i > j$ .

In the above diagram  $\mu = (8, 5, 4, 2, 1)$  and  $\alpha = \{2, 4\}$ . The grey boxes correspond to the terms in the expression which do not get cancelled.

It follows then, by the first part of proposition 2.4, that:

$$\begin{aligned} \prod_{j \in \alpha} \prod_{k=1}^{\mu_j-1} \frac{H_{\mu_-(\alpha)}(j, k)}{H_\mu(j, k+1)} &= \prod_{\substack{j \in \alpha \\ i \notin \alpha \\ i > j}} \frac{H_{\mu_-(\alpha)}(j, \mu_i)}{H_\mu(j, \mu_i+1)} \\ &= \prod_{\substack{j \in \alpha \\ i \notin \alpha \\ i > j}} \Phi(a_i, a_j)_{\epsilon q, t} \end{aligned}$$

It remains to observe that in the first column we have:

$$\prod_{j \in \alpha} \frac{1}{H_\mu(j, 1)} = \prod_{j \in \alpha} \frac{1 - a_j}{1 + ta_j/q} = w(1/t : A_J)_{\epsilon q, t}$$

□

**Proposition 2.6.** *If  $\lambda = \mu_+(\alpha, p)$  then:*

$$\begin{aligned} L(\lambda, \mu) &= \frac{(-q; t)_p}{(t; t)_p} \times \\ &\quad w(1/t : A_I)_{\epsilon q, t} V(1/t : t^{p+1} A_I)_{\epsilon q, t} W(1/t : t^p A_J)_{\epsilon q, t} \Phi(A_J, A_I)_{\epsilon q, t} \end{aligned}$$

*Proof.* The proof is similar to the previous one, only we must be slightly more careful with the first column. Begin by splitting, into two pieces, the expression:

$$\prod_{s \in \tilde{R}_{\lambda/\mu} \cap C_{\lambda/\mu}} \frac{H_\lambda(s)}{H_\mu(s)}$$

The first piece is the contribution from the first column:

$$\prod_{\substack{i=1 \\ i \notin \alpha}}^n \frac{H_{\mu_+(\alpha, p)}(i, 1)}{H_\mu(i, 1)}$$

The second piece is everything else:

$$\begin{aligned} \prod_{\substack{i \notin \alpha \\ j \in \alpha \\ i < j}} \frac{H_{\mu_+(\alpha, p)}(i, \mu_j + 1)}{H_\mu(i, \mu_j + 1)} &= \prod_{\substack{i \notin \alpha \\ j \in \alpha \\ i < j}} \frac{H_{\mu_+(\alpha)}(i, \mu_j + 1)}{H_\mu(i, \mu_j + 1)} \\ &= \prod_{\substack{i \notin \alpha \\ j \in \alpha \\ i < j}} \Phi(a_j, a_i)_{\epsilon q, t} \end{aligned}$$

Here we have made use of equation (19) to simplify the expression. We can ignore the extra “tail” of length  $p$  because it only affects the first column.

Similarly, we split, into two pieces, the expression:

$$\prod_{s \in R_{\lambda/\mu}} \frac{\tilde{H}_\lambda(s)}{\tilde{H}_\mu(s)}$$

Again the first piece is the contribution from the first column:


$$\prod_{l=1}^p \tilde{H}_{\mu_+(\alpha)}(m+l, 1) \prod_{j \in \alpha} \tilde{H}_{\mu_+(\alpha, p)}(j, 1)$$

The second piece is everything else:

$$\prod_{j \in \alpha} \prod_{k=1}^{\mu_j} \frac{\tilde{H}_{\mu_+(\alpha, p)}(j, k+1)}{\tilde{H}_\mu(j, k)} = \prod_{j \in \alpha} \prod_{k=1}^{\mu_j} \frac{\tilde{H}_{\mu_+(\alpha)}(j, k+1)}{\tilde{H}_\mu(j, k)}$$

Again we can ignore the extra “tail” of length  $p$  because it only affects the first column.

Suppose that this time we take the diagram for  $\mu$  and place it on the top of the diagram for  $\lambda = \mu_+(\alpha)$  but shifted to the right by one column.



For  $j \in \alpha$ , we have that  $H_{\mu_+(\alpha)}(j, k+1) = H_\mu(j, k)$  unless  $k = 1$  or  $k = \mu_i$  for some  $i \notin \alpha$  with  $i > j$ .



In the above diagram  $\mu = (8, 5, 4, 2, 1)$  and  $\alpha = \{2, 4\}$ . Again the grey boxes correspond to the terms which do not get cancelled.

We have, by the second part of proposition 2.4, that:

$$\begin{aligned} \prod_{j \in \alpha} \prod_{k=1}^{\mu_j} \frac{\tilde{H}_{\mu_+(\alpha)}(j, k+1)}{\tilde{H}_\mu(j, k)} &= \prod_{\substack{j \in \alpha \\ i \notin \alpha \\ i > j}} \frac{\tilde{H}_{\mu_+(\alpha)}(j, \mu_i + 1)}{\tilde{H}_\mu(j, \mu_i)} \\ &= \prod_{\substack{j \in \alpha \\ i \notin \alpha \\ i > j}} \Phi(a_j, a_i)_{\epsilon q, t} \end{aligned}$$

It remains to find suitable expressions for the four remaining factors coming from the first column:

$$\begin{aligned} \prod_{l=1}^p \tilde{H}_{\mu_+(\alpha, p)}(m+l, 1) &= \Omega((t - \epsilon q)(1 + t + \cdots + t^{p-1})) = \frac{(-q; t)_p}{(t; t)_p} \\ \prod_{j \in \alpha} \tilde{H}_{\mu_+(\alpha, p)}(j, 1) &= \prod_{j \in \alpha} \frac{1 + qt^p a_j}{1 - t^{p+1} a_j} = W(1/t : t^p A_J)_{\epsilon q, t} \\ \prod_{\substack{i=1 \\ i \notin \alpha}}^m H_{\mu_+(\alpha, p)}(i, 1) &= \prod_{\substack{i=1 \\ i \notin \alpha}}^m \frac{1 + t/q a_i t^p}{1 - a_i t^p} = V(1/t : t^{p-1} A_I)_{\epsilon q, t} \\ \prod_{\substack{i=1 \\ i \notin \alpha}}^m \frac{1}{H_\mu(i, 1)} &= \prod_{\substack{i=1 \\ i \notin \alpha}}^m \frac{1 - a_i}{1 + t/q a_i} = w(1/t : A_I)_{\epsilon q, t} \end{aligned}$$

□

**Proposition 2.7.** *The Kawanaka identity is equivalent to the following:*

$$\begin{aligned} \sum_{s=0}^k \frac{(q; t)_s}{(t; t)_s} \sum_{\substack{I \cup J = [m] \\ |J| = k-s}} &\left( w(1/t : A_I) V(1/t : t^{s-1} A_I) W(1/t : t^s A_J) \Phi(A_J, A_I) \right. \\ &\left. - w(1/t : A_J) \Phi(A_I, A_J) \right) = 0 \end{aligned}$$

*Proof.* Equation (16) may be rewritten in the form:

$$\sum_{\lambda \in \tilde{U}_k(\mu)} L(\lambda, \mu) - \sum_{s=0}^k \frac{(-q; t)_s}{(t; t)_s} \sum_{\gamma \in \tilde{D}_{k-s}(\mu)} R(\mu, \gamma) = 0$$

If  $\mu$  is a partition with distinct parts, then, as remarked at the end of the proof of proposition 2.2, we may replace the sum over vertical strips with a sum over subsets of the rows, to obtain the equivalent expression:

$$\sum_{s=0}^k \sum_{|\alpha|=k-s} \left( L(\mu_+(\alpha, s), \mu) - \frac{(-q; t)_s}{(t; t)_s} R(\mu, \mu_-(\alpha)) \right) = 0$$

The result now follows, in this special case, by applying propositions 2.5 and 2.6 and then replacing  $\epsilon q$  with  $q$  (formal negative).

If  $\mu$  contains a repeated part, and  $\alpha$  is such that  $\mu_-(\alpha)$  is not a partition then we have  $a_{i-1} = ta_i$  for some  $j = i - 1 \in \alpha$  and  $i \notin \alpha$  and so the expression:

$$\Phi(A_I, A_J) = \frac{(a_i + a_j t/q)(a_i - a_j/t)}{(a_i - a_j)(a_i + a_j/q)}$$

vanishes.

Similarly if  $\mu$  contains a repeated part, and  $\alpha$  is such that  $\mu_+(\alpha, p)$  is not a partition then we have  $1/ta_i = a_{i+1}$  for some  $i \notin \alpha$  and  $j = i + 1 \in \alpha$  and the expression:

$$\Phi(A_J, A_I) = \prod_{\substack{i \notin \alpha \\ j \in \alpha}} \frac{(a_j + a_i t/q)(a_j - a_i/t)}{(a_j - a_i)(a_j + a_i/q)}$$

vanishes.

Thus, even when  $\mu$  contains a repeated part, we may still replace the sum over vertical strips with a sum over subsets of the rows, since the terms which do not correspond to partitions all vanish. The result follows.  $\square$

The above identity is none other than a special case of proposition 2.1 with  $z = 1/t$ , which completes the proof of the Kawanaka identity:

$$\sum_{\lambda} \prod_{s \in \lambda} \left( \frac{1 + q^{a_{\lambda}(s)} t^{l_{\lambda}(s)+1}}{1 - q^{a_{\lambda}(s)+1} t^{l_{\lambda}(s)}} \right) P_{\lambda}(X; q^2, t^2) = \prod_{i \geq 1} \frac{(-tx_i; q)_{\infty}}{(x_i; q)_{\infty}} \prod_{i < j} \frac{(t^2 x_i x_j; q^2)_{\infty}}{(x_i x_j; q^2)_{\infty}}$$

## REFERENCES

- [1] F. Bergeron, *Combinatorial species and tree-like structures*, Cambridge University Press, 1998.
- [2] ———, *Combinatoire algébrique*, (2001),  
<http://bergeron.math.uqam.ca/AlgebraicCombinatorics/notes.pdf>.
- [3] Douglas Bowman, *q-difference operators, orthogonal polynomials, and symmetric expansions*, Memoirs of the American Mathematical Society Number 757 (2002).
- [4] W.Y.C. Chen, *Compositional calculus*, Journal of Combinatorial Theory, Series A **64** (1993), 149–188.
- [5] ———, *The theory of compositionals*, Discrete Mathematics **122** (1993), 59 – 87.
- [6] A. Di Bucchianico and D. Loeb, *A selected survey of the umbral calculus*, Elec. J. Combin **3** (1995).
- [7] D. Foata and G.N. Han, *The q-series in combinatorics; permutation statistics (preliminary version)*, (2004),  
<http://www-irma.u-strasbg.fr/~foata/paper/qseriescom.pdf>.
- [8] W. Fulton, *Young tableaux*, Cambridge University Press, 1997.
- [9] A.M. Garsia, *An exposé of the Mullin–Rota theory of polynomials of binomial type*, Linear and Multilinear Algebra **1** (1973), 47 – 65.
- [10] A.M. Garsia, M. Haiman, and G. M. Tesler, *Explicit plethystic formulas for Macdonald  $q, t$ -kostka coefficients*, Séminaire Lotharingien de Combinatoire **42** (1999), B42m,  
<http://www.mat.univie.ac.at/~slc/wpapers/s42garsia.html>.
- [11] A.M. Garsia and S.A. Joni, *A new expression for umbral operators and power series inversion*, Proceedings of the American Mathematical Society volume 64 (1977), 179–185.
- [12] ———, *Composition sequences*, Communications in Algebra (1980), 1187–1215.
- [13] A.M. Garsia and G. Tesler, *Plethystic formulas for Macdonald  $q, t$ -Kostka coefficients*, Advances in Mathematics **123** (1996), no. 2, 144–222.
- [14] G. Gasper and M. Rahman, *Basic hypergeometric series*, Cambridge University Press, 2004.
- [15] I.M. Gessel, *Applications of the classical umbral calculus*, Universalis **49**, 8. (2003), 397–434.
- [16] J. Goldman and G. C Rota, *On the foundations of combinatorial theory V: Eulerian differential operators*, Studies in Applied Mathematics (1971).
- [17] K. Hoffman and R. Kunze, *Linear algebra. second edition*, Prentice Hill, 1971.
- [18] E. Ihrig and M.E.H. Ismail, *A q-umbral calculus*, J. Math. Anal. Appl. **84** (1981), 178–207.
- [19] S.A. Joni and G.C. Rota, *Coalgebras and bialgebras in combinatorics*, Studies in Applied Mathematics **61** (1979), 93.
- [20] V. Kac and P. Cheung, *Quantum calculus*, New York, Springer, 2002.
- [21] N. Kawanaka, *A q-series identity involving Schur functions and related topics*, Osaka. J. Math **36** (1999), 157–176.
- [22] D. Knutson, *Lambda rings and the representation theory of the symmetric group*, Lecture Notes in Mathematics, 308 (1973).

- [23] J.P.S. Kung, *Gian-Carlo Rota on combinatorics: Introductory papers and commentaries*, Birkhäuser Boston, 1995.
- [24] A. Lascoux, *Symmetric functions and combinatorial operators on polynomials*, Regional Conference Series in Mathematics Number 99, 2001.
- [25] D. Loeb and G.-C. Rota, *Recent contributions to the calculus of finite differences: a survey*, Geometry and complex variables (Bologna, 1988/1990), Lecture Notes in Pure and Appl. Math., vol. 132, Dekker, New York, 1991, pp. 239–276, [arXiv:math/9502210](#).
- [26] D.E. Loeb, *Sequences of symmetric functions of binomial type*, Stud. Appl. Math. **83** (1990), no. 1, 1–30, [arXiv:math/9502221](#).
- [27] J. Zeng M. Ishikawa, F. Jouhet, *A generalization of Kawanaka's conjecture for Hall-Littlewood polynomials and applications*, Journal Algebraic. Combin. No 4 **23** (2006), 395–412.
- [28] I.G. Macdonald, *A new class of symmetric functions*, Séminaire Lotharingien **20** (1988), B20a, <http://www.mat.univie.ac.at/~slc/opapers/s20macdonald.html>.
- [29] ———, *Symmetric functions and Hall polynomials. second edition*, Oxford University Press, 1995.
- [30] L. Manivel, *Symmetric functions, Schubert polynomials and degeneracy loci*, SMF/AMS Texts and Monographs Volume 6, 1998.
- [31] M. Mendez, *Multiset and the combinatorics of symmetric functions*, Advances in Mathematics **102** (1993), 95–175.
- [32] M.A. Mendez, *The umbral calculus of symmetric functions*, Advances in Mathematics **124** (1996), 207 – 271.
- [33] R. Mullin and G.C. Rota, *On the foundations of combinatorial theory III: Theory of binomial enumeration*, Graph Theory and its Applications B (1970), 167–213.
- [34] K. Purbhoo, *Puzzles, tableaux, and mosaics*, (2007), [arXiv:math/0705.1184v1](#).
- [35] O. Warnaar R. Langer, M.J. Schlosser, *Theta functions, elliptic hypergeometric series and Kawanaka's macdonald polynomial conjecture*, to appear...
- [36] A. Ram, *Quantum groups: A survey of definitions, motivations and results*, (1997), <http://www.math.wisc.edu/~ram/pub/1997AustMSLectNotesv11p20.pdf>.
- [37] S. Roman, *More on the umbral calculus, with emphasis on  $q$ -umbral calculus*, Journal of Mathematical Analysis and Applications **107** (1985), 222 – 254.
- [38] ———, *The umbral calculus*, Academic Press, 2004.
- [39] G.C. Rota, *Hopf algebra methods in combinatorics*, Reprinted in his selected works.
- [40] ———, *Finite operator calculus*, Academic Press, New York, 1975.
- [41] B. Sagan, *The symmetric group*, Springer, 2001.
- [42] R. Stanley, *Enumerative combinatorics volume 2*, Cambridge University Press, 1997.
- [43] M. Zabrocki,  *$q$ -analogs of symmetric function operators*, Discrete Mathematics, Volume 256, Number 3 (2002), 831–853, [arXiv:math/0008188v2](#).
- [44] P. Zinn-Justin, *Littlewood–Richardson coefficients and integrable tilings*, Electron. J. Combin. **16** (2009), Research Paper 12.